arXiv:math/0605435v1 [math.AG] 16 May 2006

NORMAL PROJECTIVITY OF COMPLETE SYMMETRIC VARIETIES

ALESSANDRO RUZZI

Abstract

Chirivì and Maffei [CM II] have proved that the multiplication of sections of any two ample spherical line bundles on the wonderful symmetric variety $X = \overline{G/H}$ is surjective. We have proved two criterions that allows ourselves to reduce the same problem on a (smooth) complete symmetric variety to the corresponding problem on the complete toric variety (respectively to the open toric variety). We have also studied in details some family of complete symmetric varieties, in particular those of rank 2.

In this work we will study the projective normality of complete symmetric varieties. Let $\overline{G}$ be an adjoint semisimple group over $\mathbf{C}$ and let θ be an involution of $\overline{G}$. We define $\overline{H}$ as the subgroup of the elements fixed by θ and we will say that $\overline{G}/\overline{H}$ is a homogeneous symmetric variety. De Concini and Procesi [CSV I] have defined a wonderful completion of $\overline{G}/\overline{H}$ and this is the unique wonderful completion of $\overline{G}/\overline{H}$. In this work we will define a complete symmetric variety as an $\overline{G}$-variety with a dense open orbit isomorphic to $\overline{G}/\overline{H}$ and a $\overline{G}$-equivariant map $Y \rightarrow X$ extending the identity of $\overline{G}/\overline{H}$. They are be classified by De Concini and Procesi [CSV II]. Indeed they showed that there is an equivalence of categories between the category of complete symmetric varieties and the category of toric varieties over an affine space $\mathbf{A}^l$ considered as a $(\mathbf{C}^*)^l$ variety in the obvious way, where l is the rank of G/H. Moreover there is a one-to-one correspondence between the completions Y of $\overline{G}/\overline{H}$ which lie over X and the elements of a special class of complete toric varieties. One can show that the complete toric variety Z^c corresponding to a complete symmetric variety Y is a subvariety of Y and the open toric variety Z corresponding to Y is an open subvariety of Z^c.

In this work, unless explicitly stated, we shall always assume that the complete symmetric variety Y is smooth. Recall that by [CSV II] it then follows that: 1) any orbit closure in Y is also smooth; 2) the associated toric varieties Z and Z^c are both smooth.

Our work consist of two parts. In the first one we will reduce the study of the projective normality of the complete symmetric varieties to the study of projective normality of the corresponding complete toric varieties. First of all we will prove that a complete symmetric variety is projective if and only if the corresponding complete toric variety is projective. Moreover they are projective if and only if the associated open toric variety is quasi-projective.

Chirivì and Maffei [CM II] have proved the following result which easily implies the projective normality of X with respect to any projective embedding by a complete linear system.

Theorem 0.1 *Let L_1 and L_2 be any two line bundles generated by global sections on the wonderful complete symmetric variety X. Then the product of sections*

$$H^0(X, L_1) \otimes H^0(X, L_1) \longrightarrow H^0(X, L_1 \otimes L_2)$$

is surjective.

We will try to generalize this results to any complete symmetric variety. First we will prove that the surjectivity of the product of sections of two ample line bundles on a complete symmetric variety is equivalent to the surjectivity of the product of sections of the restrictions of the line bundles to the corresponding complete toric variety. Thus we will have reduced the problem to a problem on toric varieties. Unfortunately, it is very difficult to verify the surjectivity of the product of sections of any two ample line bundles on a generic complete toric variety. However, we can simplify the problem for the special class of complete toric varieties which we are considering. Indeed we will prove that the surjectivity of the product of sections of two ample line bundle on Z^c, say L_1 and L_2, is equivalent to the surjectivity of the product of sections of the restrictions of the line bundles to Z. This problem is much simpler, because $H^0(Z, L_1|Z)$ and $H^0(Z, L_2|Z)$ are infinite dimensional vector spaces and it is sufficient to prove that a suitable finite dimensional subspace of $H^0(Z, (L_1 \otimes L_2)|Z)$ is contained in the image of the product of sections. Indeed we will prove that, given any ample line bundle L on Z^c, $H^0(Z, L|Z)$ is generated by $H^0(Z^c, L|Z^c)$ as an $\mathcal{O}_Z(Z)$-module.

In the second part of this work we will study the suriectivity of the product of sections of an ample line bundle on a toric variety proper over $\mathbf{A}^l$. If the variety has dimension 2, we will prove the product of sections of any two ample line bundles is surjective. If the dimension of the variety is larger than 2, we will find a infinite number of varieties proper over $\mathbf{A}^l$ such that, for any ample line bundle L on a such variety Z, the product of sections of L

$$H^0(Z, L) \otimes H^0(Z, L) \longrightarrow H^0(Z, L \otimes L)$$

is surjective.

We would like to thank C. De Concini for the continuous help. Moreover we would like to thank W. Fulton for some useful information.

1 Introduction

Let G be a semisimple simply connected algebraic group over $\mathbf{C}$ and let θ be an involution of G. We define H as the normalizer $N_G(G^\theta)$ of the subgroup of θ-fixpoints. Let $\overline{G}$ be the adjoint semisimple group associated to G and let $\overline{H}$ be the subgroup of the elements fixed by the involution induced by θ, then $\overline{G}/\overline{H}$ is isomorphic to G/H through the map induced by the quotient map $G \to \overline{G}$.

Definition 1.1 *We will say that G/H is a homogeneous symmetric variety (of adjoint type).*

We can associate a possibly not reduced root system to G/H (see [He]). Let T^1 be a torus of G such that: 1) $\theta(t) = t^{-1}$ for each $t \in T^1$; 2) the dimension l of T^1 is maximal. Let T be a maximal torus which contains T^1. One can show that T is stabilized by θ, so θ induces an involution on $\chi^*(T)_{\mathbf{R}}$, which we call again θ. This involution stabilizes the root system ϕ of G and it is orthogonal with respect to the Killing form. We can choose a Borel subgroup B of G such that the associated set of positive roots ϕ^+ has the following property: for each $\alpha \in \phi^+$ either $\theta(\alpha)$ is equal to α or $\theta(\alpha)$ is a negative root. For each root α we define $\alpha^s = \alpha - \theta(\alpha)$. The set $\{\alpha^s \neq 0 : \alpha \in \phi\}$ is a possibly not reduced root system of rank l called the restricted root system. We will say that the not-zero α^s are restricted roots and that l is the rank of G/H. The restricted roots generate the (-1)-eigenspace of $\chi^*(T)_{\mathbf{R}}$.

We now want to describe the lattice of integral weights of ϕ^s. Let Λ be the lattice of integral weights of ϕ and let Λ^+ be the set of dominant weights. For each $\alpha \in \Gamma$, let ω_α be the fundamental weight associated to α. Moreover, for each dominant weight λ, let V_λ be the irreducible representation of G of highest weight λ. We will say that a dominant weight λ is spherical if V_λ contains a not-zero vector fixed by the Lie algebra $\mathfrak{h}$ of H. Moreover we will say that a weight μ is special if $\theta(\mu) = -\mu$. Let Ω be the lattice generated by the spherical weights and let Λ^1 be the lattice of the special weights. One can easily show that $2\Lambda^1 \subset \Omega \subset \Lambda^1$. We can describe Ω more explicitly. Let ϕ_0 be the set of root fixed by θ and let $\phi_1 = \phi - \phi_0$. We set $\Gamma_0 = \Gamma \cap \phi_0$ and $\Gamma_1 = \Gamma \cap \phi_1$ where Γ is the basis of ϕ associated to ϕ^+. The set $\Gamma^s = \{\alpha^s : \alpha \in \Gamma_1\}$ is a base of ϕ^s. θ induces an involution $\overline{\theta}$ of Γ_1 such that, for each $\alpha \in \Gamma_1$, $\theta(\alpha) = -\overline{\theta}(\alpha) - \beta_\alpha$ where β_α is a linear combination with positive integral coefficients of simple roots fixed by θ. We can order the simple roots $\alpha_1, ..., \alpha_l, \alpha_{l+1}, ..., \alpha_{l+s}, \alpha_{l+s+1}, ..., \alpha_m$ so that α_i is fixed by θ if and only if $i > l + s$. Moreover we can suppose that $\alpha_1^s, ..., \alpha_l^s$ are distinct. For each $i = 1, ...l$ we define ω_i as follows: if $\overline{\theta}(\alpha) = \alpha$ then $\omega_i = \omega_{\alpha_i}$, otherwise $\omega_i = \omega_{\alpha_i} - \theta(\omega_{\alpha_i}) = \omega_{\alpha_i} - \omega_{\overline{\theta}(\alpha_i)}$. A weight is special if and only if it is a linear combination of the weight $\omega_1, ..., \omega_l$, so $\{\omega_1, ..., \omega_l\}$ is a basis of Λ^1. Moreover we can use these weights to give the following explicit description of Ω.

Proposition 1.1 (Lemma 2.1 and Theorem 2.3 in [CM I]) *Let Ω^+ be the set of spherical weights, then $\Omega \cap \Lambda^+ = \Omega^+$. Moreover $\Omega = \bigoplus_{i=1}^{l} \boldsymbol{Z} a_i \omega_i$ where $a_i \in \{1, 2\}$ for each i. a_i is equal to 2 if $\theta(\alpha_i) = -\alpha_i$, while it is equal to 1 if $\theta(\alpha_i) \neq -\alpha_i$. In particular $a_i = 1$ if $\overline{\theta}(\alpha_i) \neq \alpha_i$. For each i and j we have $< a_i\omega_i, (\alpha_j^s)^\vee >= b_i \delta_{i,j}$ where $(\alpha_j^s)^\vee$ is the coroot associated to α_j^s and $b_i \in \{1, 2\}$. $b_j = 2$ if and only if $2\alpha_j^s \in \widetilde{\phi}$. In particular, if $\widetilde{\phi}$ is reduced then $a_1\omega_1, ..., a_l\omega_l$ are the fundamental weights dual to $(\alpha_1^s)^\vee, ..., (\alpha_l^s)^\vee$.*

Notice that the proposition implies that the fundamental Weyl chamber C^+ of $\widetilde{\phi}$ is the intersection of $M_{\mathbf{R}}$ with the fundamental Weyl chamber Λ^+ of ϕ. We will say that a special weight $\sum n_i \omega_i$ is regular if $n_i > 0$ for each i. Thus a spherical weight is regular if and only if it is a strongly dominant weight of the restricted root system (with respect to the basis Γ^s). The Weyl group W^1 of the restricted root system is called the restricted Weyl group and it has the following description.

Proposition 1.2 (See 1.8 in [CS]) *One can identify the restricted Weyl group W^1 with the group $\{w \in W : w \cdot \mathfrak{t}_1 \subset \mathfrak{t}_1\}/W_0$, where W is the Weyl group of ϕ and W_0 is the Weyl group of the root system ϕ_0 (in $\chi^*(T^0)_{\boldsymbol{R}}$).*

De Concini and Procesi have defined a wonderful compactification of G/H. Let λ be a regular spherical weight and let k_λ be a not-zero vector of V_λ fixed by $\mathfrak{h}$. One can show that k_λ is unique up to a not-zero scalar. Let x_0 be the class of k_λ in $\mathbf{P}(V_\lambda)$. De Concini and Procesi have defined the wonderful compactification X of G/H as the closure of Gx_0 in $\mathbf{P}(V_\lambda)$.

We now want to give a local description of X. Let T^0 be the connected component of the subgroup of the invariant of T and let S be the quotient of T^1

by $T^1 \cap T^0$. Choose a basis of weight vectors and consider the affine open set $\widetilde{A}$ of $\mathbf{P}(V)$ where the coordinate corresponding to the highest weight v_λ is not zero. Observe that $\widetilde{A} \cap X$ is U^- stable, where U^- is the unipotent group associated to $-\phi_1^+$, namely $U^- = \prod_{-\alpha \in \phi_1^+} U_\alpha$ as a variety. One can show that the closure of $T[k]$ in $\widetilde{A}$ is an affine space $\mathbf{A}^l$ with coordinates $-\alpha_1^s, ..., -\alpha_l^s$. Moreover the map $\varphi : U^- \times \mathbf{A}^l \to \widetilde{A} \cap X$ given by $\varphi(g, v) = g \cdot v$ is an isomorphism. For each i, let $X_{-\alpha_i^s}$ be the divisor of X whose intersection with $U^- \times \mathbf{A}^l$ is the locus of zeroes of $-\alpha_i^s$. De Concini and Procesi have proved that X is wonderful according to the definition of Luna:

Theorem 1.1 (Theorem 3.1 in [CSV I]) *Let X be as before, then:*

1. *the stabilizer of x_0 is H;*
2. *X is a smooth projective G-variety;*
3. *$X \backslash (G \cdot [k])$ is a divisor with normal crossings. It has irreducible components $X_{-\alpha_1^s}, ..., X_{-\alpha_l^s}$ and they are smooth subvarieties of X;*
4. *the G-orbits of X correspond to the subsets of $\{1, 2, ..., l\}$, so that the orbit closures are the intersections $X_{-\alpha_{i_1}^s} \cap ... \cap X_{-\alpha_{i_k}^s}$;*
5. *there is an unique closed orbit $\bigcap_{i=1}^{l} X_{-\alpha_i^s}$ and it is isomorphic to G/P, where P is the parabolic subgroup of G associated to Γ_0, i.e. its Lie algebra is $\mathfrak{t} \oplus \bigoplus_{\alpha \in \phi_0 \cup \phi^+} \mathfrak{g}_\alpha$;*
6. *X does not depend by λ.*

De Concini and Procesi have also classified the embeddings of G/H over Y i.e. the G varieties Y with a open orbit isomorphic to G/H and an equivariant map $\pi : Y \to X$ extending the identity map of G/H. We will say that a such variety is a complete symmetric variety if it is complete.

Theorem 1.2 *There is an equivalence between the category of complete symmetric varieties and the category of toric varieties proper over $\boldsymbol{A}^l$. Given a complete symmetric variety $\pi : Y \to X$ the corresponding open toric variety Z is the inverse image of $\boldsymbol{A}^l$ in Y. Moreover Y is smooth if and only if Z is smooth.*

One can show that the inverse image of the open set $U^- \times \mathbf{A}^l$ is isomorphic to $U^- \times \pi^{-1}(\mathbf{A}^l)$ in a $U^- \cdot T$ equivariant way. Moreover the G orbits of Y are in one-to-one correspondence to the S orbits of Z. The linear map $\chi^*(S)_{\mathbf{R}} \longrightarrow \chi^*(T^1)_{\mathbf{R}}$ induced in an obvious way by the quotient map $T^1 \twoheadrightarrow S$ is an isomorphism, while the linear map $\chi^*(T)_{\mathbf{R}} \longrightarrow \chi^*(S)_{\mathbf{R}}$ induced by the canonical injection $T^1 \hookrightarrow T$ is surjective. Thus we obtain a surjective map $\chi^*(T)_{\mathbf{R}} \twoheadrightarrow \chi^*(S)_{\mathbf{R}}$, whose restriction to the (-1)-eigenspace is an isomorphism. This map allow ourselves to identify $\chi^*(S)$ with the lattice M generated by the restricted roots. We call N the dual group of M, i.e. $N = Hom(M, \mathbf{Z})$.

De Concini and Procesi have also proved that there is an one-to-one correspondence between the complete symmetric varieties and a class of complete toric varieties. The closure Z_0^c of $\mathbf{A}^l$ in X is the complete toric variety whose fan is composed by the Weyl chambers and their faces.

Theorem 1.3 *There is an one to one correspondence between complete symmetric varieties and complete toric S-varieties over Z_0^c whose fan is W^1 invariant.*

Given a complete symmetric variety $\pi : Y \to X$ the corresponding complete toric variety Z^c is the inverse image of Z_0^c in Y. One can show that Z^c is the closure of Z in Y. Moreover Y is smooth if and only if Z^c is smooth. *In this work, unless explicitly stated, we shall always assume that the complete symmetric variety Y is smooth.* In this case any orbit closure in Y is also smooth.

We now want to study the line bundles on a complete symmetric variety. We begin with the wonderful case. First of all, we can identify the Picard group of the wonderful symmetric variety X with a subgroup Λ_X of the lattice Λ of integral weights of G. Indeed De Concini and Procesi have proved that:

Proposition 1.3 (Proposition 8.1 in [CSV I]) *The map $Pic(X) \to Pic(G/P)$ induced by the canonical inclusion is injective.*

Remember that we can identify $Pic(G/P)$ with a sublattice of the lattice of weights. $Pic(G/P) \equiv Pic_G(G/P)$ because G/P is a spherical variety. Thus to any linearized line bundle $L \in Pic_G(G/P)$ we associate the opposite λ of the character $-\lambda$ with which T acts on the fibre over $P \in G/P$. Because of the previous proposition, we denote a line bundle on X with L_λ if its image is the weight λ. Let λ be a dominant weight such that $\mathbf{P}(V_\lambda)$ contains a line r fixed by H, for example $\lambda \in \Omega^+$. One can show that the map $G/H \ni gH \to gr$ can be extended to a morphism $\psi_\lambda : X \to \mathbf{P}(V_\lambda)$. The line bundle $\psi_\lambda^* O(1)$ is L_λ. Thus, given a line bundle L_μ on X such that μ is dominant, there is a sub-representation of $H^0(X, L_\mu)$ isomorphic to V_μ^* and obtained by pullback of $H^0(\mathbf{P}(V_\mu), \mathcal{O}(1))$ to X. $H^0(X, L_\mu)$ is multiplicity free because X is a spherical variety, so the previous representation is unique and we can call it V_μ^*. Before to give an explicit description of $Pic(X)$ we need a definition.

Definition 1.2 *We will say that a root $\alpha \in \Gamma_1$ is an exceptional root if $\overline{\theta}(\alpha) \neq \alpha$ and $< \alpha, \theta(\alpha) > \neq 0$. Moreover we will say that G/H is exceptional if there is an exceptional root. We will say that a compactification of G/H is exceptional if G/H is exceptional.*

Theorem 1.4 (Theorem 4.8 in [CS]) *$Pic(X)$ is generated by the spherical weights and by the fundamental weights corresponding to the exceptional roots.*

We now consider the general case following [Bi]. First of all we introduce some notations that we will often use. Let X be the wonderful complete variety and let Y be the complete symmetric variety over X associated to a toric variety Z proper over $\mathbf{A}^l$. We will call Δ the fan of Z and Δ^c the fan of Z^c. We shall denote the fan of $Z_0 := \mathbf{A}^l$ by Δ_0 and the fan of Z_0^c by Δ_0^c. Let o_γ be the S-orbit of Z associated to $\gamma \in \Delta$. We will call O_τ the G-orbit of Y corresponding to o_τ. We shall denote by Z_γ the stable subvariety of Z associated to $\gamma \in \Delta$, by Z_γ^c the stable subvariety of Z^c associated to $\gamma \in \Delta^c$ and by Y_γ the stable subvariety of Y associated to $\gamma \in \Delta$. We set $\Delta(i) = \{\gamma \in \Delta : \dim \gamma = i\}$ and $\Delta^c(i) = \{\gamma \in \Delta^c : \dim \gamma = i\}$.

Remember that the closed orbits O_σ of Y are in one-to-one correspondence with the maximal cones of the fan Δ associated to Z. Moreover they are all isomorphic to the unique closed orbit of X through the restriction of the projection, so we can identify $Pic(O_\sigma)$ with Λ_X for each $\sigma \in \Delta(l)$. One can easily show that $Pic(Z)$ is freely generated by the line bundles $O(Z_\tau)$ where τ varies in the set $\Delta(1)\backslash\Delta_0(1)$. The following theorem gives a complete description of $Pic(Y)$. Given a cone $\gamma \subset N_{\mathbf{R}}$ we will call $\gamma^\perp$ the subspace of $M_{\mathbf{R}}$ of the vectors which vanishes on γ.

Theorem 1.5 (Theorem 2.4 in [Bi]) *Let Y be the complete symmetric variety associated to Z. Then*

1. *The maps* $Z \xhookrightarrow{i} Y \xrightarrow{\pi} X$ *induce the split exact sequence*
$$\begin{tikzcd} 0 \arrow[r] & Pic(X) \arrow[r, "\pi^*"] & Pic(Y) \arrow[r, "i^*"] & Pic(Z) \arrow[r] & 0, \end{tikzcd}$$
so $Pic(Y)$ is (not canonically) isomorphic to $Pic(X) \oplus Pic(Z)$.

2. *A section of the previous split short exact sequence is given by sending the free generators $\mathcal{O}(Z_\tau)$, with $\tau \in \Delta(1)\backslash\Delta_0(1)$, to $\mathcal{O}(Y_\tau)$. Thus*
$$Pic(Y) = \pi^* Pic(X) \oplus \bigoplus_{\tau \in \Delta(1)\backslash\Delta_0(1)} \boldsymbol{Z}\mathcal{O}(Y_\tau).$$

3. *The morphism given by the restriction to the closed orbits*
$$c_1^G : Pic(Y) \to \prod_{\sigma \in \Delta(l)} Pic(O_\sigma)$$
is injective and its image can be identified with the lattice
$$\Lambda_Y = \{h = (h|\sigma) \in \prod_{\sigma \in \Delta(l)} \Lambda_X \subset \prod_{\sigma \in \Delta(l)} \Lambda : \; h|\sigma - h|\sigma' \in M \cap (\sigma \cap \sigma')^\perp$$
$$\forall\ \sigma,\ \sigma' \in \Delta(l).\}$$

We will indicate with L_h the line bundle whose image is h. $Pic(Y)$ is isomorphic to the group of equivariant line bundles $Pic_G(Y)$ because G is simply connected and Y is complete. Moreover, given a line bundle L_h, $-h_\sigma$ is the character of the action of T on the fibre over the T-stable point $O_\sigma \cap Z$. In a similar way, we define h^c as the set $(h|\sigma)$ where σ varies in $\Delta^c(l)$ and $-h_\sigma$ is the character of the action of T on the fibre over the T-stable point of Z^c corresponding to σ. In some case we can give an useful interpretation of h and h^c. Before doing it, we need to give some definitions. Let M' be a lattice in $M_{\mathbf{R}}$ which contains M.

Definition 1.3 *A real valued function* $h : |\Delta| \to \boldsymbol{R}$ *on the support of* Δ *is called a* (Δ, M')*-linear function if* h *is linear on each* $\sigma \in \Delta$*. Let* $h|\sigma$ *be the unique linear function which coincide with* h *on* σ*. We request moreover that* $h|\sigma$ *belongs to* M' *and that* $h|\sigma_1 - h|\sigma_2$ *belongs to* M *for each* σ*,* σ_1 *and* σ_2 *in* $\Delta(l)$*. Let* $SF(\Delta, M')$ *be the additive group of the* (M', Δ)*-linear functions.*

Sometimes we say that an element h of $SF(\Delta, M')$ is a Δ linear function. We can think a Δ-linear function as a function $h : N_{\mathbf{R}} \to \mathbf{R} \cup \{-\infty\}$. We say that a Δ linear function h is convex if $h(v+v') \geq h(v) + h(v')$ for each $v, v' \in |\Delta|$. We say that h is strictly convex on Δ if moreover $h|\sigma \neq h|\sigma'$ for each $\sigma, \sigma' \in \Delta(l)$. Observe that, given any $h \in SF(\Delta, M')$, there is a positive integer n such that nh is (Δ, M)-linear function. One can show that, if a torus S' is a etale cover of S with character group M', then we can identify $SF(\Delta, M')$ with the group of S'-linearized line bundles on Z (S' acts on Z by the quotient map $S' \twoheadrightarrow S$). We can suppose that $-h|\sigma$ is the character of S' on the fibre $L(x_\sigma)$ over the S-stable point x_σ associated to any maximal cone σ of Δ.

Definition 1.4 *Let* L_h *be a line bundle on* Y*. We will say that* L_h *is almost spherical if* $h|\sigma$ *belongs to lattice generated by the spherical weights for each* $\sigma \in \Delta(l)$*. Moreover we will say that* L_h *is spherical if* $h|\sigma$ *is a spherical weight for each* $\sigma \in \Delta(l)$*. We will say that h is almost spherical (respectively spherical) if* L_h *is almost spherical (respectively spherical).*

Observe that if L_h is an almost spherical line bundle, then we can think h as a (Δ, Λ_X) linear function and h^c as a (Δ^c, Λ_X) linear function. We now want to describe the sections of a line bundle over Y. Observe that the space of sections is multiplicity-free because Y is a spherical variety, i.e. it has a dense B-orbit. Now we want to define sets in bijective correspondence with the sets of the irreducible subrepresentations respectively of $H^0(Y, L)$, $H^0(Z, L|Z)$ and $H^0(Z^c, L|Z^c)$.

Definition 1.5 *Given* $h \in \Lambda_Y$ *let*

$$\Pi(Z, h) = \{\mu \in \bigcap_{\sigma \in \Delta(l)} (h|\sigma + (M \cap \check{\sigma}))\},$$

$$\Pi(Z^c, h) = \{\mu \in \bigcap_{\sigma \in \Delta^c(l)} (h|\sigma + (M \cap \check{\sigma}))$$

and

$$\Pi(Y, h) = \Pi(Z, h) \cap \Lambda^+.$$

Before we describe the sections of L_h, we want to rewrite the conditions for a weight to belong to $\Pi(Z, h)$, respectively to $\Pi(Z^c, h)$.

Lemma 1.1 *Let* λ *be a weight in* Λ_X *and let* h *be in* Λ_Y*. Then:*

1. $\lambda \in \Pi(Z, h)$ *if and only if* $\lambda \geq h$ *as functions on* $|\Delta|$
2. $\lambda \in \Pi(Z^c, h)$ *if and only if* $\lambda \geq h^c$ *as functions on* $N_{\boldsymbol{R}}$

Theorem 1.6 (Theorem 3.4 in [Bi]) *Let L_h be a line bundle on Y. Then*

$$H^0(Y, L_h) = \bigoplus_{\mu \in \Pi(Y,h)} V_\mu^*.$$

In particular $H^0(Y, L_h) \neq 0$ if and only if $\Pi(Y, h)$ is not empty.

We want to give an idea of a possible construction of $H^0(Y, L_h)$. We need the following lemma.

Lemma 1.2 (Lemma 2.7 in [Bi]) *$O(Y_\tau)$ is an almost spherical line bundle whose associated Δ-linear function d^τ satisfies the equalities $d^\tau(\varrho(\tau')) = -\delta_{\tau,\tau'}$ for each $\tau' \in \Delta(1)$. Moreover, if define $s_\tau \in H(Y, L_{d^\tau})$ as the unique, up to a scalar, section $s_\tau \in H(Y, L_{d^\tau})$ whose divisor is Y_τ, then it is G invariant.*

Observe that, given $\lambda \in \Lambda_X$, then $\lambda \in \Pi(Z, h)$ if and only if $h = \lambda + \sum_{\tau \in \Delta(1)} a_\tau d^\tau$ where a_τ is a positive integer for each $\tau \in \Delta(1)$. Let $\lambda \in \Pi(Y, h)$ and remember that $H^0(X, L_\lambda)$ contains V_λ^*. $H^0(X, L_\lambda) \subset H^0(Y, \pi^* L_\lambda)$, so $H^0(Y, \pi^* L_\lambda)$ contains a lowest weight vector $v_{-\lambda}$ of weight $-\lambda$. There are positive constants a_τ such that $h - \lambda = \sum_{\tau \in \Delta(1)} a_\tau d^\tau$ because of the lemma 1.1. Thus $v_{-\lambda} \cdot \prod s_\tau^{a_\tau}$ is a not-zero section of $H^0(Y, L_h)$ with weight $-\lambda$ because the sections s_τ are G-invariant. Moreover $v_{-\lambda} \cdot \prod s_\tau^{a_\tau}$ is invariant by the unipotent part of the opposite B^- of the fixed Borel group of G. Thus $H^0(Y, L_h) \supseteq \bigoplus_{\mu \in \Pi(Y,h)} V_\mu^*$. Because of the previous theorem we give the following definition:

Definition 1.6 *Given h in Λ_Y and λ in $\Pi(Y, h)$, we write $h = \lambda + \sum_{\tau \in \Delta(1)} a_\tau d^\tau$ for suitable $a_\tau \in \boldsymbol{Z}^+$. We define $s^{h-\lambda}$ as the section $\prod s_\tau^{a_\tau}$ of $H^0(Y, L_{h-\lambda})$.*

We want to describe also the the sections over Z, respectively over Z^c.

Proposition 1.4 *Let L_h be a line bundle on Y. Then*

1. $$H^0(Z, L_h|Z) = \bigoplus_{\mu \in \Pi(Z,h)} \boldsymbol{C}\chi^\mu,$$

   where χ^μ is a T-seminvariant section of weight $-\mu$. In particular $H^0(Z, L_h|Z) \neq 0$ if and only if $\Pi(Z, h)$ is not empty.

2. $$H^0(Z^c, L_h|Z^c) = \bigoplus_{\mu \in \Pi(Z^c,h)} \boldsymbol{C}\chi^\mu.$$

   In particular $H^0(Z^c, L_h|Z^c) \neq 0$ if and only if $\Pi(Z^c, h)$ is not empty.

Remark. Let $\pi : Y \to Y'$ be an G-equivariant morphism between two complete symmetric varieties and let L_h be a line bundle on Y'. Then the pullback $\pi^*(L_h)$ is the line bundle on Y associated to h and $H^0(Y, \pi^*(L_h)) = H^0(Y', L_h)$.

Now we want to explain some relations between the previous sets.

Corollary 1.1 (Corollary 4.1 in [Bi]) *Given $h \in \Gamma_Y$, we have the equality*

$$\Pi(Y, h) = \Pi(Z^c, h) \cap \Lambda^+.$$

If L_h is almost spherical, we can say more.

Proposition 1.5 (Proposition 4.2 and theorem 4.2 in [Bi]) *If $h \in \Lambda_Y$ is almost spherical, then*

$$\Pi(Z^c, h) = \bigcup_{w \in W^1} w \cdot \Pi(Y, h).$$

Moreover the restriction map $H^0(Y, L_h) \to H^0(Z^c, L_h|Z^c)$ is surjective.

We want to make some remark on the second part of the proposition. Let $w \cdot \mu \in \Pi(Z^c, h)$ with $\mu \in \Pi(Y, h)$. Any section $s \in V_\mu^* \subset H^0(Y, L_h)$ of weight $-w \cdot \mu$ has not-zero restriction to Z^c because $U^- \cdot Z^c$ is dense in Y. Moreover, up to choose another basis of the root system, we can suppose that it is lowest weight vector.

Remember that there is an one-to-one correspondence between the convex functions on $N_{\mathbf{R}}$ with values in $\mathbf{R} \cup \infty$ and the convex sets in $M_{\mathbf{R}}$, which send a convex function h to the convex set $Q_h = \{m \in M_{\mathbf{R}} : m(v) \geq h(v),\ \forall\ v \in N_{\mathbf{R}}\}$. Moreover $h(v) = inf\{m(v);\ m \in Q_h\}$ for each $v \in N_{\mathbf{R}}$. To every linearized line bundle L_h on a toric variety we associated the polyhedron Q_h. We want to do the same with the line bundles on a complete symmetric variety. (Remember that a polyhedron is the intersection of a finite number of semispaces).

Definition 1.7 *Let Y be a complete symmetric variety and let L_h be an almost spherical line bundle on Y. We define the polytope associated to L_h as*

$$P_h = \{m \in M_{\boldsymbol{R}} : m(v) \geq h^c(v)\ \forall v \in |\Delta^c|\}.$$

Moreover we define the polyhedron associated to h as

$$Q_h = \{m \in M_{\boldsymbol{R}} : m(v) \geq h(v)\ \forall v \in |\Delta|\}.$$

Observe that $P_h = \{m \in M_{\mathbf{R}} : m(\varrho(\tau)) \geq h^c(\varrho(\tau))\ \forall\ \tau \in \Delta^c(1)\}$ and $Q_h = \{m \in M_{\mathbf{R}} : m(\varrho(\tau)) \geq h(\varrho(\tau))\ \forall \tau \in \Delta(1)\}$.

2 Ample line bundles and line bundles generated by global sections

Brion [Br] has found a characterization of the ample line bundles (respectively the line bundles generated by global sections) on a spherical variety. Now we want to find different conditions for a line bundle on a complete symmetric variety to be generated by global sections, respectively to be ample.

Proposition 2.1 *Let L_h be a line bundle on Y. Then*

1. *L_h is generated by global sections if and only if h is convex and $h|\sigma$ is dominant for each $\sigma \in \Delta(l)$.*
2. *L_h is very ample if and only if h is strictly convex on Δ and $h|\sigma$ is a regular weight for each $\sigma \in \Delta(l)$.*
3. *L_h is ample if and only if it is very ample.*

Proof. One can easily show the necessity of the conditions. Indeed if L is generated by global sections (respectively is ample) then the restrictions of L to Z and to any closed orbit O_σ are generated by global sections (respectively are ample). To prove the sufficiency of the condition in the first point we need a lemma.

Lemma 2.1 *If h is convex and $h|\sigma$ is dominant then the restriction map to the closed orbit O_σ*

$$H^0(Y, L_h) \to H^0(O_\sigma, L_h|O_\sigma)$$

is surjective.

Proof. We have $H^0(O_\sigma, L_h|O_\sigma) = V^*_{h|\sigma}$ because $h|\sigma$ is dominant. Moreover there is a lowest weight vector $\varphi \in H^0(Y, L_h)$ of weight $-h|\sigma$ because of the convexity of h. Hence, because of the irreducibility of $V^*_{h|\sigma}$, it is sufficient to prove that the restriction of φ to O_σ is not zero. Observe that $\varphi = \varphi' \cdot \prod s^{a_i}_{\tau_i}$, where φ' is a lowest weight vector of $V^*_{\lambda_\sigma} \subset H^0(Y, L_{h|\sigma})$ and $a_i > 0$ only if τ_i is not contained in σ. Recall the s_{τ_i} vanishes exactly on Y_{τ_i} and observe that $\varphi'|O_\sigma \neq 0$ because $L_{h|\sigma}|O_\sigma$ is the line bundle corresponding to $h|\sigma$. $\square$

Now we can prove the sufficiency of the condition in the first point. If the locus of base points is not empty, then it contains a closed orbit O_σ, because it is closed and stabilized by G. Given any $y \in O_\sigma$ there is a section $\widetilde{s} \in H^0(O_\sigma, L_h|O_\sigma)$ such that $\widetilde{s}(y) \neq 0$ because $h|\sigma$ is dominant. This section can be extended to a global section over Y because of the previous lemma. This is a contradiction.

Now we want to show the sufficiency of the condition in the second point if L_h is spherical. First of all we want to show that $L_h|Z$ is very ample, or better still that $L_h|Z^c$ is very ample. h^c is convex because of the first point of the proposition. Suppose by contradiction that there are two distinct maximal cones σ and σ' such that $h|\sigma = h|\sigma'$. Let γ be the convex cone formed by the points v such that $h(v) = (h|\sigma)(v)$. We can suppose that $\sigma \in \Delta$ and that $\sigma' \notin \Delta$, so there is an hyperplane H secant to γ and whose intersection with $\sigma(e_1, ..., e_l)$ is a face of $\sigma(e_1, ..., e_l)$, say $\sigma(e_1, ..., \widehat{e}_i, ..., e_l)$. Moreover there is a vector v of γ contained in the interior of $|\Delta|$ and such that also $s_i v$ belongs to γ.

By hypothesis we know that $h^c(v) = (h|\sigma)(v)$ and that $h^c(s_i v) = (s_i \cdot (h|\sigma))(v)$, so $(h|\sigma)(v) = (s_i(h|\sigma))(v)$ because of the invariance of h^c by W^1. $((h|\sigma) - s_i(h|\sigma))(v)$ is strictly positive because $h|\sigma$ is a regular weight and v is in the interior of $|\Delta|$; this is a contradiction. Observe that we have proved a more general statement. Let Z be a possibly singular toric variety proper over $\mathbf{A}^l$ and let L_{h^c} be a linearized line bundle on Z^c such that h^c is invariant for the action of W^1. If, for each $\sigma \in \Delta(l)$, $h^c|\sigma$ is a regular weight (respectively a dominant weight) and h is strictly convex (respectively convex) on the fan of Z, then L_{h^c} is ample (respectively generated by global sections).

Since L_h is generated by global sections, we have an equivariant morphism $\varphi : Y \to \mathbf{P}(V)$ with $V = H^0(Y, L)^*$. Let U be the locus where φ is not an embedding. We could try to prove this point like the previous one, namely using the fact that the restriction of L_h to Z, respectively to any closed orbit is very ample. Instead we will use the stronger fact that $L_h|Z^c$ is ample. Observe that the restriction of the sections from Y to Z^c is surjective, while the restriction of the sections to Z is clearly not surjective.

Now we want to show that U is stable and closed in the Euclidean topology. U is the union of two loci: the locus U_1 of the points where the differential of φ is not injective and the locus U_2 of the points where φ is not injective. U_1 and U_2 are G-stable because φ is equivariant. U_1 is closed because it is the locus of the zeroes of the jacobian of φ. Now we want to prove that the closure of U_2 is contained in U. Let $\{x_n\}$ be any sequence in U_2 and suppose that it converges to $x \in Y$. By hypothesis there is a sequence $\{y_n\}$ in U_2 such that $x_n \neq y_n$ and $\varphi(x_n) = \varphi(y_n)$ for each n. We can suppose, up to taking sub-sequences, that $\{y_n\}$ has limit y in Y and that $\varphi(x) = \varphi(y)$. Suppose by contradiction that x does not belong to U, in particular $x = y$. Because of the Dini theorem there is a open neighborhood W of x such that $\varphi|W$ is a diffeomorphism onto the image $\varphi(W)$. This is a contradiction because there is an integer n_0 such that x_n and y_n belong to W for each $n > n_0$. In particular, we have proved that if U_1 is empty then x must be different from y.

First suppose that U_1 is not empty, so it contains a closed orbit O_σ. Let x_σ be the intersection of Z and O_σ. The map $H^0(Z^c, L_h|Z^c)^* \longrightarrow H^0(Y, L_h)^*$ dual to the restriction map is injective, so we have a commutative diagram

$$\begin{tikzcd}
Y \arrow[r, "\varphi"] & \mathbf{P}(H^0(Y, L_h)^*) \\
Z^c \arrow[u] \arrow[r, "\varphi'"] & \mathbf{P}(H^0(Z^c, L_h|Z^c)^*) \arrow[u]
\end{tikzcd} .$$

φ' is an embedding because h^c is strictly convex on Δ^c, so $\varphi(Z^c)$ is isomorphic to Z^c. Let $[h]$ be the image $\varphi(H)$ of $H \in G/H$ and let $[v_{h|\sigma}]$ be the image of x_σ. Observe that $[v_{h|\sigma}]$ is the class of a highest weight vector of $V_{h|\sigma} \subset H^0(Y, L_h)^*$. We can write $H^0(Y, L_h)^* = V_{h|\sigma} \oplus V'$ for a suitable representation V'. We can choose $h = v_{h|\sigma} + \sum v_i$ where the v_i are weight vectors with weights contained in $h|\sigma + M$. Let $\widetilde{A}$ be the affine open set of $\mathbf{P}(H^0(Y, L_h)^*)$ where a lowest weight vector $s \in H^0(Y, L_h)$ of weight $-h|\sigma$ is not zero, i.e. $\widetilde{A} = v_{h|\sigma} + V'_{h|\sigma} \oplus V'$ where $V_{h|\sigma} = \mathbf{C}v_{h|\sigma} \oplus V'_{h|\sigma}$ in a T-equivariant way. $A := \widetilde{A} \cap \varphi(Y)$ is B^- stable and its intersection with $\varphi(Z^c)$ is $\varphi(U_\sigma)$, so $\varphi(x_\sigma)$ belongs to A. Indeed the set of points $\{x \in Z^c : \ s(x) = 0\}$ is the union of the divisor Z^c_τ for $\tau \nsubseteq \sigma$.

We want to study the restriction of φ to $U^- \cdot U_\sigma$, where U^- is the unipotent group whose Lie algebra is $\bigoplus_{\alpha \in -\phi_1^+} \mathfrak{g}_\alpha$. $H^0(Y, L_h)^*$ is isomorphic to its dual in a θ linear way (see lemma 1.6 [CSV I]). Thus there is a not degenerate bilinear form $(\,,)$ on $H^0(Y, L_h)^*$ with the following properties: 1) given any two distinct irreducible components V_1 and V_2 of $H^0(Y, L_h)^*$, they are orthogonal; 2) $(gu, v) = (u, \theta(g^{-1})v)$ for each $g \in G$, $u, v \in H^0(Y, L_h)^*$. Let Υ' be the tangent space in $v_{h|\sigma}$ to the orbit $U^- \cdot v_{h|\sigma}$ and let Υ be the space generated by Υ' and $v_{h|\sigma}$. One can show that the restriction of $(\ ,\ )$ to Υ is non-degenerate, that Υ is stable under P and that the orthogonal $\Upsilon^\perp$ is stable under $\theta(P)$ (see lemma 2.4 [CSV I]). Moreover $U_\sigma \subset v_{h|\sigma} + \Upsilon^\perp$ (see lemma 2.5 in [CSV I]). Let q be the projection of $H^0(Z^c, L_h|Z^c)^*$ onto $H^0(Z^c, L_h|Z^c)^*/\Upsilon^\perp$. $U^- \subset \theta(P)$, so U^- acts on $H^0(Z^c, L_h|Z^c)^*/\Upsilon^\perp$ and the projection is equivariant. The affine hyperplane $q(\widetilde{A})$ in $H^0(Z^c, L_h|Z^c)^*/\Upsilon^\perp$ is stable by the action of U^-. One can show that the map $j : U^- \to \pi(\widetilde{A})$ defined by $j(u) = q(uv_{h|\sigma})$ is an U^- equivariant isomorphism (see lemma 2.6 in [CSV I]). Thus the tangent space to

U_σ at $v_{h|\sigma}$ is orthogonal to Υ and the differential of φ is injective in x_σ. Hence $U_1 = \emptyset$, so U_2 is equal to U and it is closed.

Now suppose that U_2 is not empty, so it contains a closed orbit O_σ. Given $x \in O_\sigma$, there is $y \neq x$ such that $\varphi(x) = \varphi(y)$. We can suppose that y belongs to a closed orbit. Indeed there is an element g of G and an one parameter subgroup γ of T such that $y' = lim_{t\to 0}\gamma(t)gy$ is a point of Z fixed by T, in particular y' belongs to a closed orbit $O_{\sigma'}$. Moreover $x' = lim_{t\to 0}\gamma(t)gx$ belongs to O_σ and $\varphi(x') = \varphi(y')$. By the previous part of the proof x' is different by y'.

The closed orbits O_σ and $O_{\sigma'}$ are different because $L|O_\sigma$ is very ample. Because of the lemma 2.1 there is a global section s, lowest weight vector of weight $-h|\sigma$, which does not vanish on O_σ, so we can suppose that $s(x) \neq 0$ up to a translation. Since s is U^- invariant and h is strictly convex on Δ, s vanishes on a G-stable divisor of Y which contains $O_{\sigma'}$. Hence we have obtained a contradiction, namely $\varphi(x) \neq \varphi(y)$.

Finally we can consider the exceptional case. First of all we want to recall some facts. Let Y be a complete exceptional symmetric variety and let X be the corresponding wonderful variety. We can chose an order of the simple roots of ϕ such that $\alpha_1, ..., \alpha_s$ are exceptional roots with the following property: $Pic(X)$ is generated by the spherical weights and by the fundamental weights $\omega_{\alpha_1}, ..., \omega_{\alpha_s}$ corresponding respectively to $\alpha_1, ..., \alpha_s$. Moreover, given an element h of Λ_Y such that $h|\sigma$ is dominant for each $\sigma \in \Delta(l)$, there are integers a_i such that $h' = h - \sum a_i \omega_{\alpha_i}$ is another element of Λ_Y and $h'|\sigma \in \Omega$ for each $\sigma \in \Delta(l)$. For each i we can suppose that a_i is positive, up to exchange α_i with $\overline{\theta}(\alpha_i)$. If L_h satisfies the hypotheses of the second point, then $L_{h'}$ is very ample and $L_{h-h'}$ is generated by global sections, so L_h is very ample.

The third point is obvious. □

Now, we can characterize the ample line bundles on Z.

Proposition 2.2 *Suppose that Z is a (possibly singular) toric variety proper over $\boldsymbol{A}^l$ and let $h \in SF(\Delta, M)$. Then L_h is ample if and only if h is strictly convex on Δ. If Z is smooth then L_h is ample if and only if it is very ample.*

Proof. Suppose that L_h is ample, then there is an integer n such that L_{nh} is very ample, in particular nh is convex. Hence L_{nh} is the pullback of a line bundle generated by global sections on a variety Z' such that Z is proper over Z' and nh is strictly convex on the fan Δ' of Z'. Let $\varphi : Z \to \mathbf{P}(V)$ be an embedding such that $L_{nh} = \varphi^*\mathcal{O}(1)$, then φ factorizes through Z' because $H^0(Z', L_h) = H^0(Z, L_h)$. Since φ is an embedding, Z' must be Z, so h is strictly convex on Δ. The viceversa is implied by the following lemma.

Lemma 2.2 *Let Z be is a (possibly singular) toric variety proper over $\boldsymbol{A}^l$ and let L be a line bundle on Z generated by global sections. Given any homogeneous symmetric variety G/H of rank l, let Z^c be the complete toric variety associated to Z and observe that $N_{H^0}(S)$ acts on Z^c. Then there is a linearized line bundle L' on Z^c generated by global sections and a $N_{H^0}(S)$ linearization of L' such that: i) the restriction of L' to Z is L; 2) for each S-fixpoint x of Z, the character of S on the fibre $L'(x)$ is a regular weight.*

Proof. We can suppose that Z is smooth up to take a resolution of singularities of Z. Given any homogeneous symmetric variety G/H of rank l, let Y be the complete symmetric variety associated to Z. Recall that there is almost spherical line bundle L_h on Y whose restriction to Z is L and let λ be a regular spherical weight. There is a positive integer n such that $(h+n\lambda)|\sigma$ is a regular weight for each $\sigma \in \Delta(l)$, so $L' = L_{h+n\lambda}$ satisfies ours requests. $\square$.

We now can conclude the proof the proposition 2.2. Observe that, up to changing the linearization of L_h, we can suppose that the line bundle L_{h^c} is as in the previous lemma. Moreover L_{h^c} is ample on Z^c and L_h is ample on Z because of the proof of the proposition 2.1. The last point of the proposition is implied by the Demazure theorem (see [De]). $\square$

Corollary 2.1 *Let Y be a complete symmetric variety. The following conditions are equivalent:*

1. *Y is projective;*
2. *Z^c is projective;*
3. *Z is quasiprojective.*

Now we want to reformulate the proposition 2.1 using h^c instead of h.

Proposition 2.3 *Let h be an almost spherical Δ-linear function, then*

1. *h^c is convex on Δ^c if and only if h is convex on Δ and $h|\sigma$ is dominant for each $\sigma \in \Delta(l)$.*
2. *h^c is strictly convex on Δ^c if and only if h is strictly convex on Δ and $h|\sigma$ is a regular weight for each $\sigma \in \Delta(l)$.*

Proof. By the proof of the proposition 2.1 it is sufficient to prove the following facts. If h^c is an almost spherical convex Δ^c linear function then h is a spherical Δ-linear function. If h^c is also strictly convex on Δ^c, then $h|\sigma$ is regular for each $\sigma \in \Delta(l)$.

Given $\sigma \in \Delta(l)$, there is an element $w \in W^1$ such that $w \cdot h|\sigma$ is a dominant weight. $h|\sigma - w \cdot h|\sigma$ has positive values on $|\Delta|$ and $w \cdot h|\sigma$ is the restriction of h^c to $w^{-1} \cdot \sigma$. Let v be a vector in the interior of σ, then $(w \cdot h|\sigma)(v) = (h|\sigma)(v)$ because of the convexity of h^c. We have $w \cdot h|\sigma = h|\sigma$ because v is a vector inside the Weyl chamber $|\Delta|$, so $h|\sigma$ is dominant. If h^c is strictly dominant on Δ^c, then $h|\sigma$ is different from $w \cdot h|\sigma$ for each $w \in W^1$, so $h|\sigma$ is regular. $\square$

3 Reduction to the complete toric variety

In the following we will always suppose that L_h is an almost spherical line bundle, unless we will explicitly say otherwise. We start to study the multiplication of sections of two line bundles on Y. First of all, we want to show that this problem is equivalent to the similar problem on the complete toric variety Z^c associated to Y. Let L_h and L_k be any two line bundles on Y generated by global sections. Let

$$M_{h,k} : H^0(Y, L_h) \otimes H^0(Y, L_k) \longrightarrow H^0(Y, L_{h+k})$$

be the product of sections on Y and let

$$m^c_{h,k} : H^0(Z^c, L_h|Z^c) \otimes H^0(Z^c, L_k|Z^c) \longrightarrow H^0(Z^c, L_{h+k}|Z^c)$$

be the product of sections of the restrictions to Z^c of these line bundles.

Theorem 3.1 *Let* L_h *and* L_k *be two almost spherical line bundle on* Y *generated by global sections. Then* $M_{h,k}$ *is surjective if and only if* $m^c_{h,k}$ *is surjective.*

Proof. The necessity of the condition is implied by the surjectivity of the restriction maps from Y to Z^c. Indeed if $i : Z^c \to Y$ is the canonic inclusion then $m^c_{h,k} \circ (i^* \otimes i^*) = i^* \circ M_{h,k}$. Now suppose that $m^c_{h,k}$ is surjective. It is sufficient to show that the image of $M_{h,k}$ contains a basis of semi-invariant sections. If h and k are linear then $M_{h,k}$ is surjective by the theorem 0.1. In general, given $\nu \in \Pi(Y, h+k)$ there are $\lambda \in \Pi(Z^c, h)$ and $\mu \in \Pi(Z^c, k)$ such that $\nu = \lambda + \mu$. Moreover there are elements w_1 and w_2 of W^1 such that $w_1 \cdot \lambda$ and $w_2 \cdot \mu$ are dominant weights. Observe that $\nu \geq w_1 \cdot \lambda + w_2 \cdot \mu$ on $|\Delta|$. Moreover $w_1 \cdot \lambda \geq h$ and $w_2 \cdot \mu \geq k$ because h^c and k^c are convex and invariant for the action of W^1. Thus $s^{h-w_1\cdot\lambda} H^0(Y, L_{w_1\cdot\lambda}) \subset H^0(Y, L_h)$ and $s^{k-w_2\cdot\mu} H^0(Y, L_{w_2\cdot\mu}) \subset H^0(Y, L_k)$. We know that $Im\, M_{w_1\cdot\lambda, w_2\cdot\mu}$ contains a lowest weight vector $\varphi \in H^0(Y, L_{w_1\cdot\lambda+w_2\cdot\mu})$ of weight $-\nu$. Thus $s^{h+k-w_1\cdot\lambda-w_2\cdot\mu}\varphi$ is contained in $s^{h+k-w_1\cdot\lambda-w_2\cdot\mu} Im\, M_{w_1\cdot\lambda,w_2\cdot\mu} \subset Im\, M_{h,k}$ and it is not zero. □

We can prove the following proposition without assuming the surjectivity of $m^c_{h,k}$. Given two convex (Λ_X, Δ) linear function, say h and k, let $\Pi(Y, h, k)$ be the set of the weights of the lowest weight vectors contained in $Im M_{h,k}$.

Proposition 3.1 $\Pi(Y, h, k)$ *is saturated with respect to the dominant order of the roots in* $\widetilde{\phi}$.

Proof. $\Pi(Y, h+k)$ is saturated because the simple restricted roots have negative values on $|\Delta|$. Given $\nu \in \Pi(Y, h, k)$ there are two weights $\lambda \in \Pi(Y, h)$ and $\mu \in \Pi(Y, k)$ such that $\nu = \lambda + \mu$. Moreover there are element w_1, w_2 in the Weyl group W^1 such that $w_1 \cdot \lambda$ and $w_2 \cdot \mu$ are dominant weights. Observe that $\nu \geq w_1\cdot\lambda + w_2\cdot\mu$ on $|\Delta|$, so $\nu \in \prod(Y, w_1\cdot\lambda + w_2\cdot\mu)$. Let ν' be a spherical weight dominated by ν, then $\nu' \in \Pi(Y, w_1\cdot\lambda + w_2\cdot\mu)$ because this set is saturated. Let φ be a lowest weight vector of weight ν'. Because of the surjectivity of $M_{w_1\cdot\lambda,w_2\cdot\mu}$ we have $\varphi \in s^{h+k-w_1\cdot\lambda-w_2\cdot\mu} Im M_{w_1\cdot\lambda,w_2\cdot\mu} \subset Im M_{h,k}$. □

4 Reduction to the open toric variety

In this section we want to show that, given two ample line bundles on Y, the product of sections on Z^c is surjective if and only if the product of sections on Z is surjective. Moreover we will study the relation between the sections of $L|Z$ and the sections of $L|Z^c$ for any ample line bundle L on Y. Before we have to define some notations. We fix a cone $\sigma \in \Delta(l)$ and we set $v_h = h|\sigma$ for each $h \in SF(\Delta, \Lambda_X)$, so $\prod(Z, h) = Q_h \cap (M + v_h)$ and $\Pi(Z^c, h) = P_h \cap (M + v_h)$. Given any one-dimensional cone τ, we set $\rho(\tau)$ as the primitive vector of τ i.e. the more little not-zero vector of $N \cap \tau$.

$\{e_1, ..., e_l\}$ is the basis of $N_{\mathbf{R}}$ dual to the basis $\{f_1, ..., f_l\}$ of $M_{\mathbf{R}}$. We have to define a second basis $\{g_1, ..., g_l\}$ of $M_{\mathbf{R}}$ because the fundamental Weyl chamber C^+ is more easily defined using the basis the fundamental weights than the basis of the simple roots. g_i is a positive multiple of $-\omega_i$, more precisely $-g_i$ is the i-th fundamental weight of the unique reduced root system contained in $\widetilde{\phi}$ which share a basis with $\widetilde{\phi}$. $g_1, ..., g_l$ generate a lattice which contains M. Let $\{\check{g}_1, ..., \check{g}_l\}$ be the dual basis of $\{g_1, ..., g_l\}$. Observe that $C^+ = \sigma(-g_1, ..., -g_l)$. Given a point p in $M_{\mathbf{R}}$ we will use the following notations: $p = \sum x_i f_i =$

$\sum \dot{x}_i g_i$, using the "normal" coordinates for the basis $\{f_1, ..., f_l\}$ and the "dotted" coordinates for the basis $\{g_1, ..., g_l\}$. (In the following figures we consider the case in which the restricted root system is of type A_2 and Z is $\mathbf{A}^2$).



Proposition 4.1 *Let L_h be an ample spherical line bundle on Y. Then $Q_h \cap C^+ = P_h \cap C^+$ and $Q_h = P_h \cap C^+ + \sigma(f_1, ..., f_l)$.*





The equations of Q_h are of the form $\sum b_i x_i \geq b$ where the b_i are positive constants. Thus Q_h is stable by translation with respect to vectors in $\sigma(f_1, ..., f_l)$, i.e. $Q_h + \bigoplus \mathbf{R}^+(f_i) \subset Q_h$. Let H_j be the hyperplane of $M_{\mathbf{R}}$ generated by $g_1, ..., \widehat{g}_j, ..., g_l$, so the intersection of H_j and C^+ is a Weyl wall. Let s_j be the orthogonal reflection with respect to H_j. Observe that, if P_h contains a point p, then it contains all the translates of p by W^1, so it contains the orthogonal projections $\frac{1}{2}(p + s_j p)$ of p to the hyperplane H_j. Moreover there is no vertex of P_h contained in H_j, because h^c is strictly convex on Δ^c.

The function h_K associated to a polyhedron K has always finite values if and only if K is compact. Moreover, there is a decomposition in convex cones of the convex set $\{n \in N_{\mathbf{R}} : h_K(n) \in \mathbf{R}\}$ such that there is an one-to-one correspondence between the cones of such decomposition and the faces of K. Hence there is an one-to-one correspondence between the 1-dimensional cones of such decomposition and the semi-spaces that define K. Given a such cone τ the associated semi-space is $\{m \in M_{\mathbf{R}} : m(\varrho(\tau)) \geq h_K(\varrho(\tau))\}$.

First of all we will show that $P_h \cap C^+ = Q_h \cap C^+$. It is sufficient to show that $Q_h \cap C^+ \cap M_{\mathbf{Q}} \subseteq P_h$ because P_h is closed. Given any $m \in Q_h \cap C^+ \cap M_{\mathbf{Q}}$ and any $\tau \in \Delta^c(1)$ we have to show that $m(\varrho(\tau)) \geq h^c(\varrho(\tau))$. Because of the symmetry of Δ^c, there are $w \in W^1$ and $\tau' \in \Delta(1)$ such that $\varrho(\tau) = w \cdot \varrho(\tau')$. Observe that $w^{-1} \cdot m - m$ is a linear combination $\sum c_i f_i$ of the f_i with positive coefficients, so $m(\varrho(\tau)) = m(w \cdot \varrho(\tau')) = (w^{-1} \cdot m)(\varrho(\tau')) = m(\varrho(\tau')) + \sum c_i f_i(\varrho(\tau')) \geq m(\varrho(\tau')) \geq h(\varrho(\tau')) = h^c(\varrho(\tau))$. We now can conclude the proof. The decomposition in cones of $N_{\mathbf{R}}$ associated to $h_{P_h \cap C^+}$ has 1-dimensional cones $\{\sigma(\check{g}_1), ..., \sigma(\check{g}_l)\} \cup \Delta(1)$. $h_{P_h \cap C^+}$ has finite values on all $N_{\mathbf{R}}$, it is equal to h on $|\Delta|$ and vanishes on the vectors $\check{g}_1, ..., \check{g}_l$. The function associated to $\sigma(f_1, ..., f_l)$ vanishes on $|\Delta|$ and has value $-\infty$ on the complementary set. Thus their sum is the function h associated to Q_h and the proposition follows by the fact that $h_Q + h_{Q'} = h_{Q+Q'}$ for each polyhedrons Q and Q'. $\square$

We can prove a stronger statement on the ”rational” points of Q_h and P_h.

Proposition 4.2 *Let L_h be an ample spherical line bundle on Y, then $Q_h \cap (v_h + M) = P_h \cap C^+ \cap (v_h + M) + \sum_{i=1}^{l} \boldsymbol{Z}^+ f_i$.*

Remark. Observe that $H^0(Z, L|Z)$ is a $O_{Z^c}(Z^c)$-module through the restriction map $O_{Z^c}(Z^c) \to O_Z(Z)$ and $H^0(Z^c, L|Z^c)$ is a $O_{Z^c}(Z^c)$-submodule of $H^0(Z, L|Z)$. This proposition implies that $H^0(Z, L|Z)$ is generated by $H^0(Z^c, L|Z^c)$ as an $O_Z(Z)$-module.

Proof. Observe that f_j is orthogonal to H_j and let $\widetilde{f_i} = \frac{1}{2}(f_i + s_i f_j)$ for each $i \neq j$. Thus $\widetilde{f_i} \in H_j$ for each $i \neq j$ and $\{\widetilde{f_i}\}_{i \neq j}$ is a basis of H_j. $-f_i$ and $-f_j$ are distinct simple restricted roots, so $\widetilde{f_i} = f_i + d_i f_j$ for a suitable positive integer d_i. We have the following easy consequence of the proposition 4.1.

Lemma 4.1 $Q_h \cap H_j = P_h \cap H_j \cap C^+ + \bigoplus_{i \neq j} \boldsymbol{R}^+ \tilde{f}_i$.

Proof. Given $p = p' + \sum r_i f_i \in Q_h \cap H_j$ with $p' \in P_h \cap C^+$ and r_i positive constants, we have $p = \frac{1}{2}(p' + s_j p') + \sum r_i \frac{1}{2}(f_j + s_j f_i)$. $\square$

Let $R_j = \{p + a f_i \mid p \in Q_h \cap H_j \text{ and } -1/2 \leq a \leq 1/2\}$. First of all we want to describe the conditions for a point $m \in M_{\mathbf{R}}$ to belong to R_j. Fixed any j, we define another basis $u_1, ..., u_l$ of $M_{\mathbf{R}}$ such that $u_j = f_j$ and $u_i = g_i$ if $i \neq j$. The conditions for a point $p = \sum y_i u_i$ to belong to $Q_h \cap H_j$ are $y_j = 0$ plus conditions of the form $\sum_{i \neq j} n_i y_i \geq n$. Thus the conditions for a point $p = \sum y_i u_i$ to belong to R_j are the inequalities of the form $\sum_{i \neq j} n_i y_i \geq n$ that define $Q_h \cap H_j$ plus the inequalities $-1/2 \leq y_j \leq 1/2$. The following fundamental lemma is the unique part of the proof in which we will use the strictly convexity of h^c. (Remember that h^c is strictly convex on Δ^c if and only if L_h is ample).

Lemma 4.2 *R_j is contained in Q for each j.*



Proof. Observe that it is sufficient to show that $P_h \cap H_j \cap C^+ + [-1/2, 1/2] f_j \subset Q_h$ because of the previous lemma. Because of the convexity of Q_h it is sufficient to show that Q_h contains the points $p' \pm (1/2) f_j$ for each vertex p' of $P_h \cap H_j$.



Observe that the vertices of $P_h \cap H_j$ are orthogonal projections to H_j of suitable vertices of P_h. Indeed let p' a vertex of $P_h \cap H_j$ and let p be the endpoint different by p' of the segment intersection of P_h with the semi-line outgoing from p' and parallel to f_j. If p is not a vertex of P_h then p is an interior point of a segment I contained in P_h. Thus p' is an interior point of the projection of I to H_j and this segment is contained in P_h by the symmetry of P_h, a contradiction.

If $q' + af_j$ with $q' \in Q_h \cap H_j$ belongs to $M + v_h$, then $s_j(q' + af_j) = q' - af_j$, so $2a \in \mathbf{Z}$. Moreover if q is a vertex of $P_h \cap H_j$, then there is a constant a such that $q + af_j$ is a vertex of P_h, so it is sufficient to show that the intersection of P_h with the line parallel to f_j and passing through any vertex of $P_h \cap H_j$ is not a point. If there is a vertex p of $P_h \cap H_j$ without such property, then p is vertex of P_h belonging to H_j, a contradiction. $\square$

Now, we can conclude the proof of the proposition 4.2 (look to the following figure). Let p be a point contained in $Q_h \cap (M + v_h)$ and suppose that $p = \sum x_i f_i = \sum \dot{x}_i g_i$. If $\dot{x}_i \leq 0$ for each i, then $p \in P_h \cap C^+$. Otherwise there is an index j such that $\dot{x}_j > 0$. We know that $p = p' + \sum a_i f_i$ where $p' \in P \cap C^+$ and the a_i are positive constants. If $\dot{x}_j \geq 2$ then $a_j \geq 1$. Thus the point $p - [a_j]f_j = p' + (a_j - [a_j])f_j + \sum_{i \neq j} a_i f_i$ belongs to $Q_h \cap (M + v_h)$ and it has j-th coordinate with respect to $\{g_1, ..., g_l\}$ strictly less than 2 ($[a_j]$ is the integral part of a_j). Moreover, this coordinate can be at most 1 because $p - [a_j]f_j$ is a weight. We can suppose that it is exactly 1, so $p - [a_j]f_j - (1/2)f_j$ belongs to $Q_h \cap H_j$ and it is the projection of $p - [a_j]f_j$ to H_j. Thus $p - [a_j]f_j$ belong to R_j, so also $p - [a_j]f_j - f_j$ belongs to R_j and its j-th coordinate with respect to $\{g_1, ..., g_l\}$ is negative. Moreover $p - (p - [a_j]f_j - f_j)$ is a linear combination of the f_i with positive integral coefficients. If there is an index k such that $p - [a_j]f_j - f_j$ has negative k-th coordinate with respect to $\{g_1, ..., g_l\}$, then we reiterate the process. The process has to end in a finite number of steps because Q_h is contained in the semi-space $\{\sum x_i \geq h(e_1 + ... + e_l)\}$. $\square$



Now we can prove the most important theorem of this work.

Theorem 4.1 *Let L_h and L_k be two spherical ample line bundles on Y. Then $m_{h,k}$ is surjective if and only if $m^c_{h,k}$ is surjective.*

One can easily show that the theorem is equivalent to the following more combinatorial statement:

$$Q_h \cap (v_h + M) + Q_k \cap (v_k + M) = Q_{h+k} \cap (v_{h+k} + M)$$

if and only if

$$P_h \cap (v_h + M) + P_k \cap (v_k + M) = P_{h+k} \cap (v_{h+k} + M).$$

Proof. The sufficiency of the condition is easy. Given a point $p \in Q_{h+k} \cap (M + v_{h+k})$ we know that $p = p' + \sum c_i f_i$ where $p' \in P_{h+k} \cap C^+ \cap (M + v_{h+k})$ and the c_i are positive integers. Moreover there are $p_h \in P_h \cap (M + v_h)$ and $p_k \in P_k \cap (M + v_k)$ such that $p' = p_h + p_k$. Thus $p = (p_h + \sum c_i f_i) + p_k$ and $p_h + \sum c_i f_i$ belongs to $Q_h \cap (M + v_h)$.

Suppose now that $Q_h \cap (v_h + M) + Q_k \cap (v_k + M) = Q_{h+k} \cap (v_{h+k} + M)$. Let $m = \sum z_i f_i = \sum \dot{z}_i g_i$ be a point in $P_{h+k} \cap (M + v_{h+k})$. We can suppose that m belongs to C^+ by the symmetry of the polytopes P_h and P_k. By hypothesis there are two points $p'_0 \in Q_h \cap (M + v_h)$ and $q'_0 \in Q_k \cap (M + v_k)$ such that $p'_0 + q'_0 = m$. First, we will show that we can choose p'_0 and q'_0 such that p'_0 belongs to P_h. Indeed we know that $p'_0 = p_0 + w$ where $p'_0 \in P_h \cap C^+ \cap (M + v_h)$ and $w \in \bigoplus \mathbf{Z}^+ f_i$, so $m = p_0 + q_0$ where $q_0 := q'_0 + w$ belongs to $Q_k \cap (M + v_k)$.

Proceeding as in the proposition 4.2, we can define a sequence of pairs of points $\{(p_i, q_i)\}_{i=0,\dots,r}$ with the following properties: 1) $p_i \in Q_h \cap (M + v_h)$ for each i; 2) $q_i \in Q_k \cap (M + v_h)$ for each i; 3) $m = p_i + q_i$ for each i; 4) (p_0, q_0) is as before; 5) $(p_{i+1}, q_{i+1}) = (p_i + f_{j_i}, q_i - f_{j_i})$ for a suitable j_i and 6) $q_r \in P_k$. Indeed we can define the $\{q_i\}$ as in the proposition 4.2 and then we set $p_i = m - q_i$. Now it is sufficient to show by induction that we can choose the indices j_i so that p_i belongs to P_h for each i. We known that $p_0 \in P_h$. Now suppose that p_n belongs to P_h by inductive hypothesis. Suppose that $p_n = \sum x_i f_i = \sum \dot{x}_i g_i$ and $q_n = \sum y_i f_i = \sum \dot{y}_i g_i$. If $q_n \in P_k$ we define $r = n$ and there is nothing to prove. Otherwise there is an index j_n such that $\dot{y}_{j_n} > 0$ and it is sufficient to prove that $p_n + f_{j_n}$ belongs to P_h. Observe that $-\dot{x}_{j_n} > 0$, so $-\dot{x}_{j_n} \geq 1$ because it is an integer. Moreover $s_{j_n} p_n = p_n - \dot{x}_{j_n} f_{j_n}$ belongs to P_h. Thus P_h contains $p_n + f_{j_n}$ because it is convex. Thus we can choose $p_{n+1} = p_n + f_{j_n}$. □

Remark. 1) The previous theorem is valid with the weaker hypotheses that h, k are convex and that $h|\sigma$, $k|\sigma$ are regular spherical weights for each $\sigma \in \Delta(l)$. Indeed these hypotheses implies that no vertex of P_h (respectively of P_k) is contained in a Weyl wall.

2) Suppose that $h = k$ is convex and that $h|\sigma$ is a regular spherical weights for each $\sigma \in \Delta(l)$. In this case one can show that $L_h|Z$ is the pullback of an ample line bundle on a possibly singular toric variety Z' over $\mathbf{A}^l$. This suggests to consider only ample line bundles.

5 Line bundles on an exceptional complete symmetric variety

Let Y be an exceptional complete symmetric variety, let Z be the associated open toric variety and let Δ be the fan of Z. Given an ample spherical line bundle L_h over Y, we know that that the multiplication $M_{h,h}$ of sections on Y is surjective if and only if the multiplication $m_{h,h}$ of sections on Z is surjective. In this section we want to generalize this fact to the not spherical line bundles. Remember that $Pic(X)$ is generated by the spherical weights and by the fundamental weights $\omega_{\alpha_1}, ..., \omega_{\alpha_s}$ corresponding to the exceptional roots $\alpha_1, ..., \alpha_s$.

Proposition 5.1 *Let $L_{h'}$ be an ample line bundle on Y such that $M_{h',h'}$ is surjective and let $a_1, ..., a_l$ be positive integers. If we define $h = h' + \sum a_i \omega_{\alpha_i}$ then the product $M_{h,h}$ of sections of L_h over Y is surjective.*

Proof. Observe that L_h is an ample bundle on Y. We will prove the proposition by induction on $\sum a_i$. $M_{h,h}$ is trivially surjective if $\sum a_i = 0$. We need a lemma on the maps $M_{h,\omega_{\alpha_i}}$.

Lemma 5.1 *Let L_h be an ample line bundle on Y and let $\omega \in \{\omega_{\alpha_1}, ..., \omega_{\alpha_s}\}$. Then $M_{h,\omega}$ is surjective.*

Proof. In the following V_λ^* is the unique subrepresentation of $H^0(Y, L_\lambda)$ which contains a lowest weight vector v_λ of weight $-\lambda$. We have $H^0(Y, L_h) = \bigoplus_{\lambda \in \Pi(Y,h)} s^{h-\lambda} V_\lambda^*$, $H^0(Y, L_{h+\omega}) = \bigoplus_{\lambda \in \Pi(Y,h)} s^{h-\lambda} V_{\omega+\lambda}^*$ and $H^0(Y, L_\omega) = V_\omega^*$. The lemma is implied by the fact that, for each $\lambda \in \Pi(Y, h)$, $M_{h,\omega}(s^{h-\lambda} v_\lambda \otimes v_\omega)$ is a lowest weight vector of weight $-\lambda - \omega$. □

We now go back to the proposition. Let j be an index such that $a_j > 0$ and define $\widetilde{h} = h - \omega_{\alpha_j}$. We have the following commutative diagram

$$\begin{tikzcd}
H^0(Y, L_{\widetilde{h}}) \otimes H^0(Y, L_{\widetilde{h}}) \otimes H^0(Y, L_{\omega_{\alpha_j}}) \otimes H^0(Y, L_{\omega_{\alpha_j}}) \arrow[r] \arrow[d, "m_1"] & H^0(Y, L_h) \otimes H^0(Y, L_h) \arrow[dd, "M_{h,h}"] \\
H^0(Y, L_{2\widetilde{h}}) \otimes H^0(Y, L_{\omega_{\alpha_j}}) \otimes H^0(Y, L_{\omega_{\alpha_j}}) \arrow[d, "m_2"] & \\
H^0(Y, L_{2\widetilde{h}+\omega_{\alpha_j}}) \otimes H^0(Y, L_{\omega_{\alpha_j}}) \arrow[r, "M_{2\widetilde{h}+w_{\alpha_j},w_{\alpha_j}}"] & H^0(Y, L_{2h}).
\end{tikzcd}$$

m_1 is surjective by induction, m_2 and $M_{2\widetilde{h}+w_{\alpha_j},w_{\alpha_j}}$ are surjective because of the previous lemma, so $M_{h,h}$ is surjective. □

Theorem 5.1 *Let L_h be an ample line bundle on Y. If $m_{h,h}$ is surjective then $M_{h,h}$ is surjective.*

Proof. We know that, up to exchange α_i with $\overline{\theta}(\alpha_i)$ for some i in $\{1, ..., l\}$, there are positive integers $a_1, ..., a_l$ such that the line bundle $L_{h'}$, with $h' = h - \sum a_i w_i$, is spherical and ample. The restriction of L_h to Z is isomorphic to the restriction of $L_{h'}$ to Z, so $m_{h',h'}$ is surjective. Thus $M_{h',h'}$ is surjective because of the theorem 4.1. Hence $M_{h,h}$ is surjective by the previous proposition. □

6 Open projectively normal toric varieties

Now we want to describe some families of open toric varieties such that, if L_h is an ample line bundle on a such variety, then the product $m_{h,h}$ of sections is surjective. One family is formed by all the varieties of dimension 2 proper over $\mathbf{A}^2$. Moreover we will find an infinite number of varieties that have such property for every given dimension. In some cases we will prove that, given any two ample line bundles L_h and L_k on a fixed variety, then the product $m_{h,k}$ is surjective. *In the following we will identify M with $\boldsymbol{Z}^l$.*

6.1 Blow-ups of $\mathbf{A}^l$

Now we study the class of varieties that are blow-ups of $\mathbf{A}^l$ along an irreducible stable closed subvariety.

Proposition 6.1 *Let Z be the blow-up of $\boldsymbol{A}^l$ along the irreducible stable closed subvariety associated to $\sigma(e_1, ..., e_r)$. Let L_h and L_k be two line bundles generated by global sections on Z, then the product of sections $m_{h,k}$ is surjective.*

The inequalities for Q_h are $z_i \geq a_i$ for each $i = 1, .., l$ and $z_1 + ... + z_r \geq b$. The inequalities for Q_k are $z_i \geq c_i$ for each $i = 1, .., l$ and $z_1 + ... + z_r \geq d$. Here a_i, b, c_i and d are suitable integers. Let m be any point in $Q_{h+k} \cap M$, then there are $\tilde{m}_1 \in Q_h$ and $\tilde{m}_2 \in Q_k$ such that $\tilde{m}_1 + \tilde{m}_2 = m$, but they may have not integral coordinates. We want to translate $\tilde{m}_1$ with respect to a "little" vector v so that $\tilde{m}_1 + v$ and $\tilde{m}_2 - v$ will belong respectively to $Q_h \cap \mathbf{Z}^l$ and to $Q_k \cap \mathbf{Z}^l$. If $\tilde{m}_1 = (x_1, ..., x_l)$ then $x_i \geq a_i$. Let $[x_i]$ be the integral part of x_i and let $\epsilon_i = -[([x_i] - x_i)]$ (ϵ_i is 0 if x_i is an integer and it is 1 otherwise). $[x_i] + \epsilon_i \geq [x_i] \geq a_i$ because the a_i are integers. Likewise, if $\tilde{m}_2 = (y_1, ..., y_l)$ then $[y_i] + \epsilon_i \geq [y_i] \geq d_i$ and $\epsilon_i = -[([y_i] - y_i)]$. If $([y_1], ..., [y_l])$ belongs to Q_k, then we define $m_1 = ([x_1] + \epsilon_1, ..., [x_l] + \epsilon_l)$ and $m_2 = ([y_1], ..., [y_l])$. Clearly these points satisfy our requests. In the same way, if $([x_1], ..., [x_l])$ belongs to Q_h, then we define $m_1 = ([x_1], ..., [x_l])$ and $m_2 = ([y_1] + \epsilon_1, ..., [y_l] + \epsilon_l)$. Thus we can suppose that $\sum_{i=1}^{r} [x_i] \leq b$ and $\sum_{i=1}^{r} [y_i] \leq d$. We define $m_1 = ([x_1] + \epsilon_1, ..., [x_s] + \epsilon_s, [x_{s+1}], ..., [x_l])$ for an index s lesser than r and such that $h(e_1 + ... + e_r) = \sum_{i=1}^{r} [x_i] + \sum_{i=1}^{s} \epsilon_i = m_1(e_1 + ... + e_r)$. There is a such s because $\sum_{i=1}^{r} [x_i] - b$ is a negative integer, $\sum_{i=1}^{r} ([x_i] + \epsilon_i) \geq b + d - \sum_{i=1}^{r} ([y_i]) \geq b$ and $\epsilon_i \in \{0, 1\}$ for each i. Moreover we define $m_2 = m - m_1$. To verify that $m_2 \in Q_k$ it is sufficient to show that $\sum_{i=1}^{s} [y_i] + \sum_{i=s+1}^{r} ([y_i] + \epsilon_i) \geq k(e_1 + ... + e_r)$. This is implied by the inequality $m_2(e_1 + ... + e_r) = (m - m_1)(e_1 + ... + e_r) \geq (h + k)(e_1 + ... + e_r) - h(e_1 + ... + e_r)$. $\square$

Now we study a similar family of varieties, but we require that the two line bundles L_h and L_k are equal.

Corollary 6.1 *Let Z be the open toric variety obtained from $\boldsymbol{A}^l$ through the sequence of blow-ups along the subvarieties associated respectively to $\sigma(e_1, e_2)$, $\sigma(e_2, e_3)$,..., $\sigma(e_{r-1}, e_r)$. Let L_h be any line bundle generated by global sections on Z, then the product of sections $m_{h,h}$ is surjective.*

Proof. The inequalities for Q_h are: $z_i \geq a_i$ for each $i = 1, .., l$ and $z_{i-1} + z_i \geq b_i$ for each $i = 2, ..., r$, where the a_i and the b_i are suitable integers. Let $m = (x_1, ..., x_l) \in Q_{2h} \cap M = 2Q_h \cap M$. Observe that $m' = (x_1/2, ..., x_l/2) \in Q_h$ and

$m'+m' = m$. We define $\epsilon_i = -[([x_i/2]-x_i/2)]$, $m_1 = m'+(\epsilon_1, 0, \epsilon_3, ..., \epsilon_s, 0, ..., 0)$ and $m_2 = m - m_1$ for a suitable s. If r is odd then we choose $s = r$, otherwise we define $s = r-1$. These points belong to $Q_h \cap M$ because $[x_{i-1}/2]+[x_i/2]+(\epsilon_{i-1}+\epsilon_i)/2 \geq b_i$ for each i. □

6.2 Open toric varieties of dimension 2 and a singular family in dimension 3

Now we consider the family of smooth toric varieties proper over $\mathbf{A}^2$.

Theorem 6.1 *Let Z be any smooth toric variety proper over $\boldsymbol{A}^2$. Let L_{h_1} and L_{h_2} be two linearized line bundles generated by global sections and suppose that h_1 and h_2 are strictly convex on the same fan, then the product of sections m_{h_1,h_2} is surjective.*

The hypotheses mean that there is a variety Z' and two ample line bundle L'_h and L'_k over Z' such that L_h is the pullback of L'_h and L_k is the pullback of L'_k. We want to remark that Z' may be singular.

Proof. Define a scalar product (,) such that $\{f_1, f_2\}$ is a orthonormal basis. In this proof, when we will say that a side L of a polytope P is orthogonal to a vector v, we will always suppose that $(p, v) \geq 0$ for each $p \in P$ (and $(x, v) = 0$ for each $x \in L$). Notice that a plane H is the locus of zeroes of $x_1e_1 + x_2e_2 \in N$ if and only if it is orthogonal to $x_1f_1 + x_2f_2$.

Let $h_3 = h_1 + h_2$ and let Δ be the fan of Z. It is obviously sufficient to prove that $Q_{h_3} \cap M = Q_{h_1} \cap M + Q_{h_2} \cap M$. We want to decompose each Q_{h_i} in more simple polyhedrons. More precisely we will decompose each Q_{h_i} in two types of polyhedrons with vertices in M: 1) cones of form $p + \sigma(f_1, f_2)$ for a suitable point p and 2) triangles. These triangles will have a particular form, indeed we require that the fan associated to any such triangle $\widetilde{T}$ has 1-dimensional cones generated respectively by $-e_1, -e_2$ and by an element of $\sigma(e_1, e_2)$.



Let $m = (x_1, x_2)$ be any point in $Q_{h_3} \cap M$. If there is a vertex p^3 of Q_{h_3} such that m is contained in the polyhedron $p^3 + \sigma(f_1, f_2)$, then there is a maximal cone $\sigma \in \Delta$ such that $p^3 = h_3|\sigma = h_1|\sigma + h_2|\sigma$ and m belongs to $(h_1|\sigma + \sigma(f_1, f_2)) + (h_2|\sigma + \sigma(f_1, f_2)) = p^3 + \sigma(f_1, f_2)$ where $h_j|\sigma + \sigma(f_1, f_2)$ is the polyhedron associated to the pullback of a line bundle on $\mathbf{A}^2$. Otherwise there are two maximal cones σ_1 and σ_2 with the following properties. For each j let $p_1^j = h_j|\sigma_1 = (z_1^{1,j}, z_2^{1,j})$, $p_2^j = h_j|\sigma_2 = (z_1^{2,j}, z_2^{2,j})$, $p_3^j = (z_1^{2,j}, z_2^{1,j})$ and define T^j as the triangle with vertices p_1^j, p_2^j and p_3^j. Then $T^3 = T^1 + T^2$ and T^3 contains m. Indeed we can choose σ_1 and σ_2 such that: 1) p_1^3, p_2^3 are the two vertices of a side of Q_{h_3}; 2) $z_1^{1,3} \leq x_1 \leq z_1^{2,3}$. Observe that $x_2 \leq z_2^{1,3}$ because otherwise m belongs to $(z_1^{1,3}, z_2^{1,3}) + \sigma(f_1, f_2)$. Moreover the fans associated to these triangles are equal to the fan with 1-dimensional cones $\sigma(-e_1), \sigma(-e_2)$ and $\sigma(a_1e_1 + a_2e_2)$ for suitable integers a_1 and a_2. Observe that p_1^j (respectively p_2^j) is the vertex of T^j not contained in the side orthogonal to $-f_1$ (respectively to $-f_2$). Moreover we can suppose that $a_1f_1 + a_2f_2 \in \sigma(f_1 + f_2, f_2)$ up to exchange f_1 and f_2. In the pictures we consider the case in which $h_1 = h_2$.

We want to prove that $(T^1 + \sigma(f_1, f_2)) \cap M + (T^2 + \sigma(f_1, f_2)) \cap M = (T^1 + T^2 + \sigma(f_1, f_2)) \cap M$ by induction on $a_1 + a_2$. If $a_1 + a_2 = 2$ the previous equality is trivially implied by the proposition 6.1. Otherwise we decompose T^j in two triangles by intersecting T^j with the line r^j orthogonal to $f_1 + f_2$ and passing for the vertex p_1^j. Observe that r^j intersects the side orthogonal to $-f_1$ in the integral point $q^j = (z_1^{2,3}, z_2^{1,3} + z_1^{1,3} - z_1^{2,3})$. We have decomposed T^j in two triangles: 1) the triangle T_1^j with sides orthogonal respectively to $-f_1$, $-f_2$, $f_1 + f_2$ and with vertices p_1^j, p_3^j, q^j; 2) the triangle T_0^j with sides orthogonal respectively to $-f_1$, $-f_1 - f_2$, $a_1f_1 + a_2f_2$ and with vertices p_1^j, q^j, p_2^j. Observe that r_i^1, r_i^2 and r_i^3 are parallel, so the fans associated to T_1^1, T_1^2 and T_1^3 (respectively to T_0^1, T_0^2 and T_0^3) are equal. $T_1^3 = T_1^1 + T_1^2$ because each T_1^j share two vertices with T^j. In the same way we obtain that $T_0^3 = T_0^1 + T_0^2$. Notice that $\sigma(f_1 + f_2, f_2) \subset \sigma(f_1, f_2)$ so $T_0^j + \sigma(f_1 + f_2, f_2) \subset T^j + \sigma(f_1, f_2)$ for each j. Moreover $a_1f_1 + a_2f_2 = (a_1 - a_2)f_1 + a_2(f_1 + f_2)$ and $a_1 < a_1 + a_2$, so we can apply the inductive hypothesis. $\square$



Now we consider a class of line bundles on varieties of dimension 3. This line bundles are the pullbacks of ample lines bundles on varieties which are usually singular.

Proposition 6.2 *Let Δ be the fan with maximal cones $\sigma(e_1, e_2, ae_1 + ae_2 + e_3)$, $\sigma(e_1, e_3, ae_1 + ae_2 + e_3)$ and $\sigma(e_2, e_3, ae_1 + ae_2 + e_3)$ where a is a strictly positive integer. If h is a strictly convex Δ-linear function, then $Q_{2h} \cap M = Q_h \cap M + Q_h \cap M$.*





Remember that h defines a line bundle generated by global sections on every toric variety proper over the toric variety associated to Δ. Moreover the toric variety associated to Δ is proper over $\mathbf{A}^3$ and it is smooth if and only if $a = 1$. (Look to the figure for an example).

Proof. We want to proceed as in the previous theorem. We again define a scalar product such that $\{f_1, f_2, f_3\}$ is an orthonormal basis and we can again decompose Q_{2h} in a simplex P_0 and some cones $p + \sigma(f_1, f_2, f_3)$. Hence we can reduce ourselves to prove by induction on a that $(2P + \sigma(v_1, v_2, v_3)) \cap M = (P + \sigma(v_1, v_2, v_3)) \cap M + (P + \sigma(v_1, v_2, v_3)) \cap M$ for each simplex P with faces orthogonal respectively to $-v_1, -v_2, -v_3$ and to $av_1 + av_2 + v_3$, where $\{v_1, v_2, v_3\}$ is an opportune base of $\mathbf{Z}^3$ and a is a positive integer (observe that the polytope P_0 satisfies such requests). We can suppose, up to a translation, that the origin 0 is the vertex of P which does not belong to the face orthogonal to $av_1 + av_2 + v_3$. Let $(-b, 0, 0)$, $(0, -b, 0)$ and $(0, 0, -c)$ be the other vertices of P. We have $c = ba$, so $c \geq b$. If $a = 1$ the equality is verified because of the proposition 6.1. Otherwise we decompose P intersecting it with the plane orthogonal to $v_1 + v_2 + v_3$ and passing through the vertices $(-b, 0, 0)$ and $(0, -b, 0)$. This plane intersects the side of P parallel to $\mathbf{R}v_3$ in $(0, 0, -b)$. We obtain two simplices with integral vertices. The first one has faces orthogonal respectively to $-v_1$, $-v_2$, $-v_3$ and to $v_1 + v_2 + v_3$. This simplex has vertices $(0, 0, 0)$, $(-b, 0, 0)$, $(0, -b, 0)$ and $(0, 0, -b)$. The second simplex has faces orthogonal respectively to $-v_1$, $-v_2$, $-v_1 - v_2 - v_3$ and to $av_1 + av_2 + v_3$. This simplex T has vertices $(-b, 0, 0)$, $(0, -b, 0)$, $(0, 0, -b)$ and $(0, 0, -c)$. Observe that $\sigma(v_1, v_2, v_1 + v_2 + v_3)$ is contained in $\sigma(v_1, v_2, v_3)$, so $T + \sigma(v_1, v_2, v_1 + v_2 + v_3)$ is contained in $P + \sigma(v_1, v_2, v_3)$. Moreover T is a simplex of the same type of P and $av_1 + av_2 + v_3 = (a - 1)v_1 + (a - 1)v_2 + (v_1 + v_2 + v_3)$, i.e. the coordinate with respect to the new basis are decreased. $\square$



6.3 Stable subvarieties

In some case we can reduce the study of the product of sections of two ample line bundles to the study of the product of sections of the restrictions of these line bundles to irreducible stable closed subvarieties. First of all, Brion has proved the following proposition.

Proposition 6.3 (See theorem 4 in [BI]) *Let L be any line bundle generated by global sections on a smooth quasi-projective toric variety Z. Suppose that either Z is complete or it is proper over $\boldsymbol{A}^l$. Then, given two cones $\gamma \subset \gamma'$ in Δ, the restriction*

$$H^0(Z_\gamma, L|Z_\gamma) \longrightarrow H^0(Z_{\gamma'}, L|Z_{\gamma'})$$

is surjective.

Proposition 6.4 *Let L_h and L_k be two ample linearizated line bundles on a smooth toric variety Z. Suppose that either Z is complete or it is proper over $\boldsymbol{A}^l$. Let τ be a cone in $\Delta(1)$ and let s be a global section of L_{h+k} which does not vanish on Z_τ. If $s|Z_\tau$ belongs to the image of the product $m^\tau_{h,k}$ of sections of the restrictions of L_h and L_k to Z_τ, then s belongs to the image of the product $m_{h,k}$ of sections of L_h and L_k.*

Proof. Since there is a basis of semi-invariant sections, we can suppose that s is a semi-invariant section of weight μ, so $\mu(\varrho(\tau)) = (h+k)(\varrho(\tau))$. Because of the previous proposition, there are sections $s'_i \in \Gamma(Z^c, L_h)$ and $s''_i \in \Gamma(Z^c, L_k)$ such that $m^\tau_{h,k}(\sum s'_i|Z^c_\tau \otimes s''_i|Z^c_\tau)$ is $s|Z^c_\tau$, so $m_{h,k}(\sum s'_i \otimes s''_i) = s$. $\square$

6.4 Two families of open toric varieties of dimension at least 3

Now we want to show that there is an infinite number of open toric varieties of any fixed dimension (greater than 2) such that the product of sections of any two ample line bundles is surjective. The principal instrument in what follows is the proposition 6.4. We will consider a very special class of varieties. Indeed, given any ample line bundle L on a variety of this family, then $H^0(Z, L)$ is generated as a $O_Z(Z)$ module by the sections that do not vanish on a suitable divisor.

Proposition 6.5 *Let Z^n be the open toric variety obtained from A^l through the sequence of blow-ups along the irreducible stable subvarieties associated respectively to $\sigma(e_1, ..., e_l)$, $\sigma(e_1, ..., e_{l-1}, (\sum_{i=1}^{l-1} e_i) + e_l)$, $\sigma(e_1, ..., e_{l-1}, 2(\sum_{i=1}^{l-1} e_i) + e_l)$,...,$\sigma(e_1, ..., e_{l-1}, i(\sum_{i=1}^{l-1} e_i) + e_l)$,...,$\sigma(e_1, ..., e_{l-1}, (n-1)(\sum_{i=1}^{l-1} e_i) + e_l)$. Let L_h and L_k be any two line bundles generated by global sections on Z^n, then the product of sections $m_{h,k}$ is surjective.*



Proof. In the figure we have drawn the variety Z^3 with dimension 3. Observe that we have already considered the case $n = 1$ in proposition 6.1, so we can suppose $n \geq 2$. Up to changing the linearizations of the line bundles we can suppose that $h(e_j) = k(e_j) = 0$ for each j. Observe that, if $(Q_h \cap M) + (Q_k \cap M)$ contains a weight p, then it contains any weight $p + \sum a_i f_i$ where the a_i are positive integers. So we can consider only the "minimal" weights. We now prove the propriety of Z stated before the proposition.

Claim 6.1 *Let p be any weight in $Q_{h+k} \cap M$ and suppose that there is not a weight p' in $Q_{h+k} \cap M$ such that $p \in p' + \sigma(f_1, ..., f_l)$. Then there is a cone $\tau \in \Delta(1)$ such that $p(\varrho(\tau)) = (h + k)(\varrho(\tau))$.*

Proof. Observe that $p(\varrho(\tau)) = (h + k)(\varrho(\tau))$ means that any semi-invariant section of weight p does not vanish on the divisor Z_τ. The hypotheses imply that $p - f_l$ does not belong to Q_{h+k}. Hence there is an i such that $(p - f_l)(i(\sum_{i=1}^{l-1} e_i) + e_l) < (h + k)(i(\sum_{i=1}^{l-1} e_i) + e_l)$, so $p(i(\sum_{i=1}^{l-1} e_i) + e_l) = (h + k)(i(\sum_{i=1}^{l-1} e_i) + e_l)$. □

Now it is sufficient to prove the surjectivity of the product of sections of the restrictions of L_h and L_k to the divisor Z_i^n associated to $\sigma(i(\sum_{i=j}^{l-1} e_j) + e_l)$ for each $i = 0, ..., n$. Z_0 is the blow-up of A^{l-1} in the stable point; Z_n^n is the projective space of dimension $l - 1$ while the other Z_i^n are isomorphic to the blow-up of projective space in a point. Since Z_1^n dominates Z_n^n it is sufficient to study the product of sections of any two line bundles generated by global sections on Z_1^n.

Lemma 6.1 *Let $L_{h'}$ and $L_{k'}$ be any two line bundles on Z_1^n generated by global sections. Then the multiplication of sections is surjective.*

Proof. We can suppose that $h'(\widetilde{e}_i) = k'(\widetilde{e}_i) = 0$ for each i. In the following we identify $\mathbf{Z}^{l-1}$ with the character group of the torus contained in Z_1^n. We proceed as in the proof of the proposition 6.1. Given any point m in $Q_{h'+k'}$ with integral coordinates, there are $\tilde{m}_1 \in Q_{h'}$ and $\tilde{m}_2 \in Q_{k'}$ such that $\tilde{m}_1 + \tilde{m}_2 = m$. Now we want to simplify the notation. In particular we will be evident that the problem does not depend on the dimension. Suppose that $\tilde{m}_1 = (x_1, ..., x_{l-1})$, $\tilde{m}_2 = (y_1, ..., y_{l-1})$ and $m = (z_1, ..., z_{l-1})$. Let $[x_i]$ be the integral part of x_i and

let $\epsilon_i = -[([x_i]-x_i)]$. We define $m_1 = ([x_1]+\epsilon_1, ..., [x_{\bar{r}}]+\epsilon_{\bar{r}}, [x_{\bar{r}+1}], ..., [x_{l-1}])$ and $m_2 = ([y_1], ..., [y_{\bar{r}}], [y_{\bar{r}+1}]+\epsilon_{\bar{r}+1}, ..., [y_{l-1}]+\epsilon_{l-1})$ for a suitable $\bar{r}$. Let $t=\sum_{i=1}^{l-1}\epsilon_i$, $r=\sum_{i=1}^{\bar{r}}\epsilon_i$, $[x]=\sum_{i=1}^{l-1}[x_i]$, $x=\sum_{i=1}^{l-1}x_i$, $[y]=\sum_{i=1}^{l-1}[y_i]$ and $y=\sum_{i=1}^{l-1}y_i$. Let $a=h'(\sum_{i=1}^{l-1}\widetilde{e}_i)$, $b=-h'(-\sum_{i=1}^{l-1}\widetilde{e}_i)$, $c=k'(\sum_{i=1}^{l-1}\widetilde{e}_i)$ and $d=-k'(-\sum_{i=1}^{l-1}\widetilde{e}_i)$. We know the following inequalities: i) $[x]\leq x\leq [x]+t$, $[y]\leq y\leq [y]+t$ and $0\leq r\leq t$; ii) $a\leq x\leq b$ and $c\leq y\leq d$; iii) $a+c\leq [x]+[y]+t=x+y\leq b+d$. Observe that $\widetilde{m}_1(\sum_{i=1}^{l-1}\widetilde{e}_i)=x$, $m_1(\sum_{i=1}^{l-1}\widetilde{e}_i)=[x]+r$, $\widetilde{m}_2(\sum_{i=1}^{l-1}\widetilde{e}_i)=y$ and $m_2(\sum_{i=1}^{l-1}\widetilde{e}_i)=[y]+t-r$. It is sufficient to show that there is r such that $0\leq r\leq t$, $a\leq [x]+r\leq b$ and $c\leq [y]+t-r\leq d$. Observe that r takes all the value between 0 and t when $\bar{r}$ varies between 0 and $l-1$.

1) If $t+[x]\leq b$ we define r as $\min\{[y]+t-c, t\}$. If $[y]\geq c$ then $r=t$, so $b\geq [x]+t=[x]+r\geq x\geq a$ and $c\leq [y]\leq y\leq d$. If $c\geq [y]$ then $b\geq [x]+t\geq [x]+r=[x]+[y]+t-c\geq a+c-c=a$ and $c=[y]+t-([y]+t-c)=[y]+t-r\leq d$.

2) Suppose now that $[y]+[x]+t\leq b+c$. If $c-[y]$ is positive then we define $r=t+[y]-c$, so $t-r=c-[y]$ ($t+[y]\geq y\geq c$ so $r\geq 0$). In this case $c=[y]+t-r\leq d$ and $a\leq [x]+[y]+t-c=[x]+r\leq b$. If $c-[y]$ is negative then we define $r=t$, so $c\leq [y]=[y]+t-r\leq d$ and $a\leq x\leq [x]+t=[x]+r\leq c+b-[y]\leq b$.

3) Finally suppose that $t+[x]>b$ and $[y]+[x]+t>b+c$. We define $r=b-[x]$, so $a\leq [x]+r=b$ and $d\geq [y]+[x]+t-b=[y]+t-r\geq c$. $\square$

We have proved that the "minimal" weights of $\prod(Z^n, h+k)$ come from semi-invariant sections that do not vanish on a suitable divisor. Moreover we do not need to assume that L_h and L_k are ample. This fact are no longer true if we consider varieties whose fans are a little less symmetric. Notice that the fans of the varieties considered in the previous proposition are invariant for any automorphism of N which permutes the vectors of the basis, fixing e_l. In the following we define a class of varieties without such symmetry and obtained by blow-ups from varieties of the previous family.

Theorem 6.2 *Let $\widetilde{Z}^n$ be the blow-up of Z^n along the stable subvariety associated to $\sigma(\sum_{j=1}^{l}e_j, e_2, ..., e_l)$. Let L_h be any ample line bundles on $\widetilde{Z}^n$, then the product of sections $m_{h,h}$ is surjective.*


e_3
$\sum_{i=1}^{l} e_i$
$e_1 + 2\sum_{j=2}^{l} e_j$
$2(\sum_{i=1}^{l-1} e_i) + e_l$
$3(\sum_{i=1}^{l-1} e_i) + e_l$
e_1
e_2

Proof. In the figure we have drawn the case in which $l=3$ and $n=3$. We introduce some notation to simplify the counts: $w := e_1 + 2\sum_{j=2}^{l} e_j$ and $v_i := i(\sum_{i=1}^{l-1} e_i) + e_l$ for each i. Moreover we suppose that $h(e_j)=0$ for each j.

In the proof we allow L_h to be the pullback of an ample linearized line bundle on Z^n, i.e. $h(w) = h(v_1)$. We want to prove the proposition by induction on $h(w)$ and on the dimension of $\widetilde{Z}^n$. Observe that if $h(w) = h(v_1)$ then $m_{h,h}$ is surjective because of the previous proposition, while if the dimension of $\widetilde{Z}^n$ is 2, then $m_{h,h}$ is surjective because of the theorem 6.1. Suppose now that $h(w) > h(v_1)$. We want to prove that $Q_{2h} \cap M = (Q_h \cap M) + (Q_h \cap M)$ in a similar way to the previous proposition. Let $a_i = h(v_i)$ and $b = h(w)$. As before, if p belongs to $(Q_h \cap M) + (Q_h \cap M)$ then $p + \bigoplus \mathbf{Z}^+ f_i$ is contained in $(Q_h \cap M) + (Q_h \cap M)$. Thus we can suppose that $m - f_l$ does not belong to $(Q_{2h} \cap M)$, so either there is an i such that $m(v_i) = a_i$ or $m(w) - 2h(w) \in \{0, 1\}$.

Either we have to study a divisor or $m(w) = 2h(w) + 1$. One can easily verify that the only divisor that we do not have already considered is $\widetilde{Z}^n_{\sigma(v_1)}$. This variety can be studied in a very similar way to the divisor Z^n_1 of Z^n, so we left the details to the reader. Thus we can suppose that $m(w) = 2b + 1$. We now want to write some necessary conditions to the strictly convexity of h on the fan Δ associated to Z. The conditions $(h|\sigma(v_1, w, e_2, ..., e_{l-1}))(v_i) > h(v_i)$, $h|\sigma(w, e_2, ..., e_l)(v_1) > h(v_1)$, $h|\sigma(v_1, e_1, e_3, ..., e_l)\ (e_2) > h(e_2)$, $h|\sigma(v_1, e_1, e_3, ..., e_l)$ $(w) > h(w)$ and $h|\sigma(v_1, e_1, e_3, ..., e_l)(v_i) > h(v_i)$ imply:

$$a_i + (i-1)b < (2i-1)a_1 \quad \forall i$$

$$b > a_1 > 0, \quad 2a_1 > b, \quad ia_1 > a_i.$$

Let Δ be the fan of $\widetilde{Z}^n$, let Δ' be the fan of Z^n and let h' be the Δ linear function such that $h'(e_i) = 0$, $h'(v_i) = h(v_i)$ and $h'(w) = h(w) - 1$. We need the following easy lemma on h' which we will not prove.

Lemma 6.2 *h' is convex on Δ and is strictly convex either on Δ or on Δ'.*

By induction we can suppose that $m_{h',h'}$ is surjective, so we can suppose that there are two points $m_1 \in Q_h \cap M$ and $m_2 \in Q_{h'} \cap M$ such that $m_1 + m_2 = m$. If m_2 belong to Q_h then $m \in Q_h \cap M + Q_h \cap M$. Otherwise we have $m_2(w) = b - 1$ and $m_1(w) = b + 2$. Write $m_1 = (x_1, ..., x_l)$ and $m_2 = (y_1, ..., y_l)$.

We can suppose that $m_1 - f_l \notin Q_h$ because $m_2 + f_l \in Q_h$. Thus there is i such that $m_1(v_i) = a_i$. Moreover we can suppose that $(m_1 + f_1 - f_j, m_2 - f_1 + f_j)$ does not belong to $Q_h \times Q_{h'}$ for any $j = 2, ..., l-1$, so $x_j = 0$ or $y_1 = 0$. If $y_1 = 0$ then $2a_1 - 1 \geq b - 1 = m_2(w) = 2\sum y_j = 2m_2(v_1) \geq 2a_1$, so we have obtained a contradiction. Hence $y_1 \neq 0$ and $x_j = 0$ for each $j = 2, ..., l-1$. Suppose that there is $i > 1$ such that $m_1(v_i) = ix_1 + x_l = a_i$, then we have

$$(2i-1)a_1 \leq (2i-1)(x_1 + x_l) = m_1(v_i) + (i-1)m_1(w) =$$

$$= a_i + (i-1)(b+2) \leq (2i-1)a_1 + 2i - 2,$$

so $0 \leq (2i-1)(x_1 + x_l - a_1) \leq 2i - 2$ (remember that $a_i + (i-1)b < (2i-1)a_1$). We have $x_1 + x_l = a_1$ because $x_1 + x_l - a_1$ is an integer. Observe that we have showed that $m_1(v_1) = x_1 + x_l = a_1$ or $m_1(e_l) = x_l = 0$. In the last case we have $x_2 = ... = x_l = 0$ and $x_1 = b + 2$. We can suppose that $(m_1 - f_1, m_2 + f_1)$ does not belong to $Q_h \times Q_{h'}$, so there is $s > 0$ such that $m_1(v_s) - a_s < s$. Observe that $m_1(v_s) = sx_1 = sb + 2s$, so $a_s \leq sb = m_1(v_s) - 2s < a_s - s < a_s$, so we have obtained a contradiction.

Finally we can suppose that $x_j = 0$ for each $j = 2, ..., l-1$, $x_1 + x_l = a_1$ and $x_1 + 2x_l = b+2$, so $x_1 = 2a_1 - b - 2$ and $x_l = b + 2 - a_1$. Moreover we can suppose that $(m_1 + f_1 - f_l, m_2 - f_1 + f_l)$ does not belong to $Q_h \times Q_{h'}$, so $x_l = 0$, $y_1 = 0$ or there is $i > 1$ such that $\varepsilon := m_2(v_i) - a_i < i$. Observe that we have already considered the first two cases.

We have $a_i \leq m_1(v_i) = ix_1 + x_l = (2i-1)a_1 - (i-1)b - 2(i-1)$, so $(2i-1)a_1 \geq a_i + (i-1)b + 2(i-1)$. Finally

$$(2i-1)a_1 \leq (2i-1)(\sum y_j) \leq (2i-1)y_1 + (3i-2)\sum_{j\neq 1,l} y_j + (2i-1)y_l =$$

$$= m_2(v_i) + (i-1)m_2(w) = (i-1)(b-1) + a_i + \varepsilon =$$

$$= (i-1)b + a_i + 2(i-1) - 3(i-1) + \varepsilon \leq (2i-1)a_1 - 3(i-1) + \varepsilon,$$

so $3(i-1) \leq \varepsilon \leq i-1$, a contradiction. □

References

[Bi] E. BIFET, On Complete Symmetric Varieties, Adv in Math. **80**, (1990), 225-249.

[Br] M. BRION, Groupe de Picard et nombres caractéristiques des variétés sphérique, Düke Math. J. **58**, (1989), no. 2,397-424.

[BI] M. BRION and S. P. INAMDAR, Frobenius splitting of spherical varieties, in: Algebraic groups and their generalizations, Proc. Symp. Pure Math. **56**, AMS, Providence 1994, pp. 207-218.

[CS] C. De CONCINI and T. A. SPRINGER, Compactifications of symmetric varieties, Transform. Groups **4** (2000), no. 2-3, 273-300.

[CM I] R. CHIRIVÌ AND A. MAFFEI, The ring of sections of a complete symmetric variety, J. Algebra, **261**, no.2, 310-326, (2003).

[CM II] R. CHIRIVÌ AND A. MAFFEI, Projective normality of complete symmetric varieties, Duke Math. J. **122** (2004) no. 1, 93-123.

[CSV I] C. De CONCINI AND C. PROCESI, Complete symmetric varieties, in Invariant Theory, "Lecture notes in Math" **996**, 1-44, Springer-Verlag, New York 1983.

[CSV II] C. De CONCINI AND C. PROCESI, Complete symmetric varieties II, *Adv. Stud. Pure Math.* **6** (1985), 481-513.

[De] M. DEMAZURE, Sousgroupes algébriques de rang maximum du groupe de Cremona, Ann. Sci. École Norm. Sup. (4) **3** (1970), 507-588.

[He] S. HELGASON, "Differential Geometry, Lie groups ans Symmetric Spaces", Academic Press, San Diego, 1978.

[O] T. ODA, "Convex bodies and algebraic geometry, An introduction to the theory of toric varities, Springer-Verlag, 1985.